

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 39/395, C07K 15/28, C12N 5/12</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/26307</b> <b>(43) International Publication Date:</b> 24 November 1994 (24.11.94)
<b>(21) International Application Number:</b> PCT/US94/05433 <b>(22) International Filing Date:</b> 16 May 1994 (16.05.94) <b>(30) Priority Data:</b> 063,399 17 May 1993 (17.05.93) US <b>(71) Applicant:</b> THE PICOWER INSTITUTE FOR MEDICAL RESEARCH [US/US]; 350 Community Drive, Manhasset, NY 11030 (US). <b>(72) Inventors:</b> BUCALA, Richard, J.; 504 East 63rd Street, #33D, New York, NY 10021 (US). MITCHELL, Robert, A.; 516 East 11th Street, #6C, New York, NY 10009 (US). BERNHAGEN, Jorgen; 1570 First Avenue, #7H, New York, NY 10028 (US). CALANDRA, Thierry, F.; 500 East 63rd Street, New York, NY 10021 (US). CERAMI, Anthony; Ram Island Drive, Shelter Island, NY 11964 (US). <b>(74) Agents:</b> CORUZZI, Laura, A. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).		<b>(81) Designated States:</b> AU, BB, BG, BR, BY, CA, CN, CZ, FL, GE, HU, JP, KG, KR, KZ, LK, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, UA, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> INHIBITION OF MIGRATION INHIBITORY FACTOR IN THE TREATMENT OF DISEASES INVOLVING CYTOKINE-MEDIATED TOXICITY		
<b>(57) Abstract</b> <p>The present invention relates to compositions and methods for inhibiting the release and/or biological activity of migration inhibitory factor (MIF). In particular, the invention relates to the uses of such compositions and methods for the treatment of various conditions involving cytokine-mediated toxicity, which include, but are not limited to shock, inflammation, graft versus host disease, and/or autoimmune diseases.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INHIBITION OF MIGRATION INHIBITORY FACTOR  
IN THE TREATMENT OF DISEASES INVOLVING  
CYTOKINE-MEDIATED TOXICITY

1. INTRODUCTION

5       The present invention relates to compositions and  
methods for inhibiting the release and/or biological  
activity of migration inhibitory factor (MIF). In  
particular, the invention relates to the uses of such  
10       compositions and methods for the treatment of various  
conditions involving cytokine-mediated toxicity, which  
include, but are not limited to shock, inflammation,  
graft versus host disease, and/or autoimmune diseases.

2. BACKGROUND OF THE INVENTION

15

2.1. CYTOKINES IN SEPTIC SHOCK

      Septic shock is a multifaceted pathological  
condition characterized most prominently by  
deleterious hemodynamic changes and coagulopathy  
20       leading to multiple organ failure and often to death.  
The altered physiological mechanisms underlying the  
septic shock syndrome, and the cellular means by which  
these changes are induced and controlled, are not yet  
known in precise detail. In broad outline, however, a  
25       consensus view of events culminating in septic shock  
has emerged over the last several years. In  
particular, it is now generally accepted that septic  
shock reflects the individual, combined and concerted  
effects of a large number of endogenous, host-derived  
30       mediator molecules. These mediators are produced in  
response to initiating stimuli that indicate the host  
has been invaded, and the class of peptide mediators  
that were generally first recognized as white blood  
cell products have come to be known as cytokines. As  
35       mediators of toxic effects and pathological

alterations in host homeostasis, these endogenous factors represent potentially attractive therapeutic targets, and septic shock remains a potentially lethal cytokine-mediated clinical complication against which  
5 there is no generally effective therapeutic approach.

Although traditionally termed "septic" shock, infection by a variety of microorganisms including not only bacteria but also viruses, fungi, and parasites can induce septic shock. In fact, the shock syndrome  
10 is more properly associated with the host's response to invasion rather than just infection, as cancer and trauma, for instance, can also serve as initiators. In the case of infection by gram-negative bacteria, one of the best studied examples, it is believed that  
15 the appearance of bacterial endotoxins such as lipopolysaccharide (LPS) in the host bloodstream leads to the endogenous production of a variety of host factors that directly and indirectly mediate the toxicity of LPS, which itself is relatively innocuous  
20 for most cells. These host-derived mediators include many now well-recognized inflammatory cytokines and classical endocrine hormones in addition to a number of other endogenous factors such as leukotrienes and platelet activating factor. It is generally  
25 acknowledged, however, that the full cast of participants and each of their interrelated roles in the host response remains incompletely appreciated.

In general, those mediators that appear earlier in an invaded host are thought to trigger the release  
30 of later appearing factors. Also, many endogenous mediators not only exert direct effector functions at their target tissues, but also prime local and remote tissues for subsequent responses to other mediators. This interacting network of host factors has been  
35 termed the "cytokine cascade." This term is meant to

indicate the rapid extension and amplification of the host response in such a way that only one or a few initiating stimuli trigger the eventual release and participation of scores of host mediators. Although a  
5 number of features of the host response are thought to assist in fighting off invasion, an overly robust or poorly modulated endogenous response can rapidly accelerate to rapidly produce such profound alterations in host homeostasis at the cellular,  
10 tissue, and systemic levels that death may ensue within hours.

Among the interacting factors that together comprise the cytokine cascade, the cytokine known as tumor necrosis factor-alpha (TNF $\alpha$ ) is the most  
15 important identified to date. TNF $\alpha$  is the first cytokine to appear in the circulation after LPS challenge. The hemodynamic and metabolic alterations that result from the experimental administration of TNF $\alpha$  closely resemble those that have been observed in  
20 endotoxemia and septic shock. In animal models, TNF $\alpha$  is the only host factor which itself can initiate a lethal syndrome that mimics septic shock in detail. In this respect, TNF $\alpha$  can be considered a sufficient mediator of septic shock. Functionally neutralizing  
25 TNF $\alpha$  antagonists such as anti-TNF $\alpha$  antibodies are protective in otherwise lethal bacterial infections, and in this respect TNF $\alpha$  can be considered a necessary mediator of septic shock.

Other cytokines participate in the host response to LPS but appear later in the circulation. However,  
30 no other cytokine has been shown to be both necessary and sufficient to mediate septic shock. For example, certain interleukins (IL-1, IL-6 and IL-8) which appear in serum more than 2 hours after LPS challenge,  
35 and interferon  $\gamma$  (IFN- $\gamma$ ) which appears after 6 hours,

are thought to play a significant role in septic shock, and can be shown to contribute to lethality in certain disease models or under experimental conditions of endotoxemia. Antagonism of the effects of specific interleukins and interferons has been shown to confer a significant protective effect under certain conditions. Nevertheless, none of these other factors can itself induce a full-blown septic shock-like effect in otherwise healthy individuals, and none of these other cytokines appears to play as central and critical role in septic shock as  $\text{TNF}\alpha$ .

In view of the foregoing,  $\text{TNF}\alpha$  stands as an ideal target for the treatment of septic shock. Unfortunately, temporal characteristics of the endogenous  $\text{TNF}\alpha$  response suggest a significant practical limitation for this potential therapy.  $\text{TNF}\alpha$ , one of the earliest elicited mediators to appear in acute disease, rapidly peaks after bolus endotoxin challenge (30-90 minutes), and diminishes just as promptly. It is presumed that most of the damaging effects of  $\text{TNF}\alpha$  in septic shock are completed during this early period, hence  $\text{TNF}\alpha$  antagonists such as anti- $\text{TNF}\alpha$  antibodies would ideally be present at this time. Since this therapeutic window is apparently so short and occurs so early, the timely delivery of anti- $\text{TNF}\alpha$  based therapeutics may be very difficult to achieve clinically.

Therefore, in order utilize cytokines as targets for the treatment of septic shock and other cytokine-mediated toxic reactions, there exists a desperate need to discover additional targets that are both necessary components of the cytokine cascade and occur at a time during the endogenous response that is accessible for therapeutic antagonism in the course of clinical treatment.

## 2.2. THE PITUITARY AS A SOURCE OF PROTECTIVE CYTOKINES

Recent studies suggest that the pituitary gland may produce factors that inhibit endotoxin-induced TNF $\alpha$  and IL-1 production, and thus may serve as a source for potentially protective factors that may be used to treat shock and/or other inflammatory responses. (Suzuki et al., 1986, Am. J. Physiol. 250: E470-E474; Sternberg et al., 1989, Proc. Natl. Acad. Sci. USA 86: 2374-2378; Zuckerman et al., 1989, Eur. J. Immunol. 19: 301-305; Edwards III et al., 1991a, Endocrinol. 128: 989-996; Edwards III et al., 1991b, Proc. Natl. Acad. Sci. USA 88: 2274-2277, Silverstein et al., 1991, J. Exp. Med. 173:357-365). In these studies, hypophysectomized mice (i.e., animals that have had their pituitary glands surgically removed) exhibited a marked increased sensitivity to LPS injection relative to sham-operated control mice. In fact, the LPS LD<sub>100</sub> for control mice was approximately 1-2 logs higher than that determined for the hypophysectomized mice, suggesting that the pituitary gland produces one or more factors that may act to increase the host animal's ability to resist endotoxin challenge. Some of these studies implicate the involvement of ACTH and adrenocorticosteroids (e.g., Edwards III et al., 1991a and 1991b, supra); however, other data suggest the existence of other protective factors derived from the pituitary.

Very recently, murine macrophage migration inhibitory factor (MIF) was identified as an LPS-induced pituitary protein (Bernhagen et al., 1993, J. Cell. Biochem. Supplement 17B, Abstract E306). While it may be hypothesized that MIF is one of such protective factors capable of counteracting the adverse effects of cytokines in endotoxaemias, its

role in septic shock had not been investigated prior to the present invention.

### 2.3. MIF: MACROPHAGE MIGRATION INHIBITORY FACTOR

5        Although MIF was first described over 25 years ago as a T cell product that inhibits the random migration of guinea pig macrophages (Bloom & Bennett, 1966, Science 158: 80-82; David, 1966, Proc. Natl. Acad. Sci. USA 65: 72-77), the precise role of MIF in  
10 either local or systemic inflammatory responses has remained largely undefined. MIF has been reported to be associated with delayed-type hypersensitivity reactions (Bloom & Bennett, 1966, supra; David, 1966, supra), to be produced by lectin-activated T-cells  
15 (Weiser et al., 1981, J. Immunol. 126: 1958-1962), and to enhance macrophage adherence, phagocytosis and tumoricidal activity (Nathan et al., 1973, J. Exp. Med. 137: 275-288; Nathan et al., 1971, J. Exp. Med. 133: 1356-1376; Churchill et al., 1975, J. Immunol. 115: 781-785). Unfortunately, many of these studies  
20 used mixed culture supernatants that were shown later to contain other cytokines such as IFN- $\gamma$  and IL-4 that also have migration inhibitory activity (McInnes & Rennick, 1988, J. Exp. Med. 167: 598-611; Thurman  
25 et al., 1985, J. Immunol. 134: 305-309).

Recombinant human MIF was originally cloned from human T cells (Weiser et al., 1989, Proc. Natl. Acad. Sci. USA 86: 7522-7526), and has been shown to  
activate blood-derived macrophages to kill  
30 intracellular parasites and tumor cells in vitro, to stimulate IL-1 $\beta$  and TNF $\alpha$  expression, and to induce nitric oxide synthesis (Weiser et al., 1991, J. Immunol. 147: 2006-2011; Pozzi et al., 1992, Cellular Immunol. 145: 372-379; Weiser et al., 1992,  
35 Proc. Natl. Acad. Sci. USA 89:8049-8052; Cunha et al.,



1993, J. Immunol. 150:1908-1912). Until very recently, however, the lack of a reliable source of purified MIF has continued to hamper investigation of the precise biological profile of this molecule.

5

### 3. SUMMARY OF THE INVENTION

The present invention relates to compositions and methods which inhibit the release and/or biological activity of migration inhibitory factor (MIF). The invention further relates to the uses of such compositions and methods for the treatment of conditions involving cytokine-mediated toxicity, which include, but are not limited to shock, inflammation, graft versus host disease, and/or autoimmune diseases.

The inhibition of MIF activity in accordance with the invention may be accomplished in a number of ways which include, but are not limited to, the use of MIF binding partners, i.e., factors that bind to MIF and neutralize its biological activity, such as neutralizing anti-MIF antibodies, soluble MIF receptors, MIF receptor fragments, and MIF receptor analogs; the use of MIF-receptor antagonists, such as anti-MIF-receptor antibodies, inactive MIF analogs that bind but do not activate the MIF-receptor, small molecules that inhibit MIF release, or alter the normal configuration of MIF, or inhibit productive MIF/MIF-receptor binding; or the use of nucleotide sequences derived from MIF gene and/or MIF receptor gene, including coding, non-coding, and/or regulatory sequences to prevent or reduce MIF expression by, for example, antisense, ribozyme, and/or triple helix approaches. Any of the foregoing methods may be utilized individually or in combination to inhibit MIF release and/or activity in the treatment of the relevant conditions. Further, such treatment(s) may

be combined with other therapies that (a) inhibit or antagonize initiators of cytokine-mediated toxicity (e.g. anti-LPS antibody); (b) inhibit or antagonize toxic participants in the endogenous cytokine  
5 responses (e.g. anti-TNF $\alpha$ , anti-IL-1, anti-IFN- $\gamma$ , or IL-1 RA); or (c) themselves inhibit or antagonize cytokine-mediated toxicity (e.g. steroids, glucocorticoids or IL-10).

The present invention is based, in part, on the  
10 surprising discoveries that, first, administration of MIF in vivo increases mortality of animals after challenge with endotoxin and, second, that inhibition of MIF activity in vivo results in enhanced or prolonged survival after either challenge with  
15 endotoxin or with TNF $\alpha$ , which challenge would otherwise result in death due to shock. Prior to the present invention, no role for MIF in the inflammation/shock response was appreciated. Moreover, the existing evidence and observations  
20 indicated, at best, that MIF might play a beneficial role in the treatment of endotoxin-induced shock. First, MIF was shown to enhance macrophage killing of intracellular parasites and tumor cells. Second, MIF is an endotoxin-induced protein expressed by the  
25 pituitary, an organ which had previously been suggested as a source of protective rather than exacerbative factors involved in the shock syndrome. In contrast to such expectations, the Applicants have discovered that MIF activity actually exacerbates  
30 endotoxin-induced shock, and that inhibition of MIF activity can be used to successfully treat otherwise lethal effects of cytokine-mediated toxicity.

The invention is illustrated by working examples which demonstrate that MIF exacerbates

35

endotoxin-induced shock, and that anti-MIF prevents endotoxin-induced lethality in animal models.

#### 4. BRIEF DESCRIPTION OF THE FIGURES

- 5 FIG. 1: Graph depicting the schematic profile of  
TNF $\alpha$  and MIF responses involved in LPS-  
induced shock.
- FIG. 2: Potential ribozyme cleavage sites of MIF.  
Murine (mmif) and human (hmif) nucleotide  
10 sequences are compared. Underlined  
sequences indicate the potential ribozyme  
cleavage sites.
- FIG. 3: Nucleotide sequence of cDNA encoding murine  
MIF.
- 15 FIG. 4: Predicted amino acid sequence homology  
between murine pituitary MIF and human  
Jurkat T-cell MIF. Potential N-linked  
glycosylation sites are underlined. mu:  
murine MIF; hu: human MIF.
- 20 FIG. 5: Inhibition of monocyte migration by  
recombinant MIF. The effect of various  
concentrations (0.0001-10  $\mu$ g/ml) of rmMIF  
on peripheral human blood monocyte migration  
was quantitated in a modified Boyden  
25 chamber. The percent migration inhibition  
relative to buffer controls is plotted  
versus the logarithm of MIF concentration.  
Each point depicts the mean of independently  
performed duplicate determinations.
- 30 FIG. 6A: MIF potentiates LPS-induced lethality in  
BALB/c mice. Results represent the pooled  
data from two experiments.
- FIG. 6B: Anti-MIF antiserum protects against LPS  
challenge. Results represent the pooled  
35 data from two experiments.

- FIG. 7: Anti-MIF antiserum protects against Gram-positive staphylococcal exotoxin challenge.
- FIG. 8: Western blot analysis of MIF released by pituitary cells.
- 5 FIG. 9A and 9B: Immunocytochemistry of MIF release by pituitary cells. FIG. 9.A. Cells cultured in the absence of LPS; FIG. 9.B. Cells cultured in the presence of 25  $\mu\text{g/ml}$  LPS.
- 10 FIG. 10: LPS-induced expression of pituitary MIF mRNA in BALB/c mice.
- FIG. 11: Competitive PCR quantitation of pituitary MIF mRNA from LPS-stimulated BALB/c mice.
- FIG. 12: Presence of MIF protein in mouse pituitaries.
- 15 FIG. 13: Temporal analysis of MIF levels, in sera of mice treated with LPS.
- FIG. 14: Western blotting and densitometric analysis of serum MIF in BALB/c, BALB/c<sup>nu/nu</sup>, and hypophysectomized BALB/c mice. Mice were injected i.p. with LPS and blood sampled at intervals for Western blotting analysis. The content of MIF in serum aliquots (5 $\mu\text{l}$ ) was quantified by laser densitometry with reference to electrophoresed rMIF standards. The inset shows the MIF blot for the BALB/c mice, demonstrating the time-dependent increase in serum MIF after LPS administration. Each plotted point is the mean $\pm$ SEM of individual sera from 2 to 5 animals.
- 20
- 25
- 30
- 35 FIG. 15: Western blotting analysis of the MIF content of non-stimulated monocytes/macrophages, T-lymphocytes, and PMNs. Ten  $\mu\text{l}$  of cell lysate (equivalent to  $5 \times 10^3$  cells) were electrophoresed through 18% gels,

transferred to nitrocellulose membrane, and analyzed with anti-MIF polyclonal antibody. 30 ng of rMIF was electrophoresed and transferred as a standard.

5 FIG.  
16A, 16B,  
16C:

Western blotting analysis of MIF secretion by RAW 264.7 cells (FIG. 16A) and by thioglycollate-elicited peritoneal macrophages stimulated with LPS (FIG. 16B) or with IFN $\gamma$  plus LPS (FIG. 16C). Macrophages ( $4 \times 10^6$  RAW 264.7 cells or  $1 \times 10^7$  peritoneal macrophages) were incubated for 12 h with LPS or IFN $\gamma$  plus LPS at the indicated concentrations. The content of MIF secreted into the medium was analyzed by Western blotting. rMIF (30 ng in panel A and 20 ng in panels B and C) was electrophoresed and transferred as a standard.

20 FIG. 17: RT-PCR analysis of MIF, TNF $\alpha$ , and  $\beta$ -actin expression by LPS-stimulated RAW 264.7 macrophages. Cells ( $4 \times 10^6$ ) were incubated for 12 h in medium containing RPMI/1% FBS and LPS at the indicated concentrations. Total cellular RNA was extracted, CDNA prepared, and gene-specific PCR products analyzed by agarose gel electrophoresis as described in *Materials and Methods*.

30 FIG. 18: Competitive PCR analysis of macrophage MIF mRNA after LPS stimulation. RAW 264.7 cells ( $4 \times 10^6$ ) were incubated for 12 h in RPMI/1% FBS with or without LPS (100 pg/ml). RNA was extracted and MIF mRNA levels analyzed as described in *Material and Methods*. The

35

amount of MIF competitor template DNA ranged from 0.25 to 15 pg per reaction as shown.

FIG. 19: Western blotting analysis of cytokine-induced MIF secretion by RAW 264.7

5 macrophages. Cells ( $4 \times 10^6$ ) were incubated for 12 h with  $\text{TNF}\alpha$ , IL- $1\beta$ , IL-6 (10 or 1 ng/ml), or IFN $\gamma$  (1000 or 100 IU/ml). 20 ng of rMIF was electrophoresed and transferred as a standard.

10 FIG. 20: Concentration of  $\text{TNF}\alpha$  in cell-culture supernatants of RAW 264.7 macrophages stimulated with rMIF. Cells ( $4 \times 10^6$ ) were incubated for 12 h with the indicated concentrations of rMIF or with medium alone  
15 (control). Culture medium then was removed and the  $\text{TNF}\alpha$  content quantified by L929 cell cytotoxicity.  $\text{TNF}\alpha$  concentration is expressed as the difference between the level produced by rMIF-stimulated cells and  
20 by control (non-stimulated) cells. Data are expressed as the mean  $\pm$  SD of three separate experiments.

FIG. 21: Western blotting analysis of the inhibition by 20 $\alpha$ -dihydrocortisol of the MIF response  
25 of RAW macrophages to dexamethasone.

FIG. 22: Western blotting analysis of the organ distribution of murine MIF protein. Organs were obtained from an untreated BALB/c mouse. Aliquots (60  $\mu\text{g}$  of protein) of  
30 liver, spleen, kidney, adrenal, brain and lung lysates were electrophoresed through 18% gels, transferred to nitrocellulose membrane, and analyzed with anti-MIF polyclonal antibody. 20 ng of rMIF was  
35 electrophoresed and transferred as a

standard. Although not visible, a MIF band of much lower intensity also was detected in the adrenals and lungs.

5           5. DETAILED DESCRIPTION OF THE INVENTION

The present invention involves compositions and methods that inhibit MIF release and/or activity in vivo, for the treatment of any conditions involving cytokine-mediated toxicity, which include but are not  
10 limited to shock, including endotoxin-induced septic shock and exotoxin-induced toxic shock, inflammation, graft versus host disease, autoimmune diseases, acute respiratory distress syndrome, granulomatous diseases, chronic infections, transplant rejection, cachexia,  
15 viral infections, parasitic infections, including for instance malaria, and/or bacterial infections.

The inhibition of MIF activity in accordance with the invention may be accomplished in a number of ways, which may include, but are not limited to, the use of  
20 factors which bind to MIF and neutralize its biological activity; the use of MIF-receptor antagonists; the use of compounds that inhibit the release of MIF from cellular sources in the body; and the use of nucleotide sequences derived from MIF  
25 coding, non-coding, and/or regulatory sequences to prevent or reduce MIF expression. Any of the foregoing may be utilized individually or in combination to inhibit MIF activity in the treatment of the relevant conditions, and further, may be  
30 combined with any other anti-cytokine therapy (including steroid therapy), anti-initiator therapy, inhibitory cytokines or any combination thereof.

5.1. MIF IS A CRITICAL MEDIATOR IN THE SHOCK  
SYNDROME AND INTEGRATES CENTRAL AND  
PERIPHERAL INFLAMMATORY RESPONSES

MIF is identified herein as both a neuroendocrine mediator and a macrophage cytokine which plays an  
5 important role in host inflammatory responses to infection and tissue invasion. Upon release after a proinflammatory stimulus, macrophage-derived MIF may act together with TNF $\alpha$  and other cytokines to  
10 coordinate local cellular responses against infection or tissue invasion. Pituitary-derived MIF, however, may serve to prime systemic immune responses once a localized inflammatory site fails to contain an  
invasive agent, or else act as a CNS-derived stress  
15 signal to activate the immune system against invasion. Thus, MIF may act in concert with adrenocorticotrophic hormone (ACTH) and the adrenocortical axis to modulate systemic inflammatory responses.

The working examples described infra, demonstrate that MIF, historically considered to be a product of  
20 activated T-lymphocytes, is also produced by the pituitary and macrophages (see Section 11 and 12; infra); that MIF plays a critical role in endotoxin-induced septic shock (caused by gram negative organisms; see Section 7, infra), exotoxin-induced  
25 toxic shock syndrome (caused by gram positive organisms; see Section 8, infra), parasite-induced disease responses (see Section 9, infra), and non-infectious inflammatory diseases, e.g., involving the cell-mediated immune response (Section 10, infra); and  
30 that MIF acts in concert with glucocorticoids to regulate inflammation and immunity (see Section 13, infra). In particular, the results show that MIF potentiates lethality of endotoxemia, whereas the inhibition of MIF confers protection against lethal  
35 endotoxemia. Inhibition of MIF similarly confers



protection against toxic shock syndrome.

Surprisingly, MIF is induced by many of the glucocorticoids that are considered to be anti-inflammatory agents.

5       The experiments in animals described herein indicate that a dominant role of MIF is to serve as a proinflammatory mediator. A proinflammatory spectrum of action for MIF has been verified by in vitro studies described herein. As shown in Section 6,  
10 infra, recombinant (rMIF) was found to induce TNF $\alpha$  secretion by macrophages, indicating that MIF and TNF $\alpha$  may act locally in a reciprocal, proinflammatory loop, i.e., MIF and TNF $\alpha$  stimulate the production of each other by cells of the immune system. Recombinant MIF,  
15 in combination with other proinflammatory stimuli, e.g., IFN- $\gamma$ , also promotes murine macrophage nitric oxide (NO) synthesis (see Section 6, infra).

The Applicants have found that the resting pituitary contains large stores of preformed MIF which  
20 are released into the circulation after activation of the hypothalamic-pituitary axis by endotoxin, but not from pituicytes by several other cytokines which participate in the proinflammatory response, such as TNF $\alpha$ , IL-1 $\beta$ , IL-6 or IFN- $\gamma$  (see Section 11, infra).  
25 MIF is also released by macrophages in response to low doses of endotoxin, and in response to TNF $\alpha$  and IFN- $\gamma$  (see Section 12, infra), in response to parasitic infection (see Section 9 infra), and in response to steroids (see Section 13, infra). Thus, the  
30 macrophage is not only a target for MIF, but is also an important source of MIF in vivo.

Figure 1 depicts the profile of endotoxin-induced MIF and TNF $\alpha$  serum concentrations detected during the shock response. Detectable serum concentrations of  
35 TNF $\alpha$  quickly peak and clear within 1-2 hours of

exposure to endotoxin. In contrast, detectable serum concentrations of MIF gradually increase during the acute phase (1-8 hours), peak at 8 hours and persist during the post-acute phase (> 8 hours) for up to 20 hours. While not limited to any theory of operation, the following cellular sources are proposed for the serum concentrations of MIF observed, based on the data presented in Sections 11 and 12 infra, and Figures 13 and 14: MIF is probably produced by activated T-cells and macrophages during the proinflammatory stage of endotoxin-induced shock, e.g., as part of the localized response to infection. Once released by a pro-inflammatory stimulus, e.g., low concentrations of LPS, or by TNF $\alpha$  and IFN- $\gamma$  (but not IL-1 $\beta$  or IFN- $\gamma$ ), macrophage-derived MIF is the probable source of MIF produced during the acute phase of endotoxic shock. Both the pituitary, which releases MIF in response to LPS (but not INF- $\alpha$ , IL-1 $\beta$ , IL-6 or IFN- $\gamma$ ), and macrophages are the probable source of MIF in the post-acute phase of endotoxic shock, when the infection is no longer confined to a localized site.

In view of Applicants' discovery that the pituitary is a source of MIF, the notion that hypophysectomy sensitizes experimental animals to the lethal effects of LPS initially suggested that MIF may act to down-regulate host inflammatory responses. Therefore, it was surprising to find that purified, recombinantly prepared MIF (rMIF), when injected in mice produced visible symptoms of endotoxemia, and when co-injected with LPS, potentiated lethality in mice. E.g., see the results presented in the working example in Section 7, infra, which demonstrate that MIF substantially amplifies endotoxin-induced mortality of animals. Thus, contrary to what was

expected, it is shown herein that an increase in MIF is not beneficial to the host system, and, in fact, appears to exacerbate cytokine-mediated conditions such as septic shock. The results also show that MIF  
5 is induced by toxic shock syndrome toxin-1 (TSST-1), a staphylococcal exotoxin (Section 8, infra), and by hemozoin, a specific malarial product (Section 9, infra).

It was equally surprising to discover that  
10 inhibition of MIF conferred protection against lethal toxemia. As demonstrated in the working example presented in Section 7, infra, inhibition of MIF activity leads to a dramatic increase in the cumulative survival of animals challenged with high-  
15 dose endotoxin or TNF $\alpha$ . Anti-MIF antiserum also markedly reduced mortality in test animals challenged with lethal doses of TSST-1 (see Section 8, infra). These data establish that MIF contributes significantly to lethality and may in this sense be  
20 considered a necessary or essential component of the shock syndrome.

In accordance with the invention, the inhibition of MIF activity and/or release may be used to treat inflammatory response and shock. Beneficial effects  
25 may be achieved by intervention at both early and late stages of the shock response.

While not limited to any theory or mechanism responsible for the protective effect of MIF inhibition, preliminary studies show that anti-MIF  
30 therapy is associated with an appreciable (up to 35-40%) reduction in circulating serum TNF $\alpha$  levels. This reduction is consistent with the TNF $\alpha$ -inducing activity of MIF on macrophages in vitro, and suggests that MIF is responsible, in part, for the extremely  
35 high peak in serum TNF $\alpha$  level that occurs 1-2 hours

after endotoxin administration despite the fact that MIF cannot be detected in the circulation at this time. Thus, MIF inhibition therapy may be beneficial at the early stages of the inflammatory response.

5 MIF also plays an important role during the post-acute stage of the shock response, and therefore, offers an opportunity to intervene at late stages where other treatments, such as anti-TNF $\alpha$  therapy, are ineffective. The working examples described herein  
10 show that MIF is released at the post-acute stage of shock, when it is detectable in the circulation. In the experimental system used, anti-MIF therapy protected against lethal shock in animals challenged with high doses of endotoxin (i.e., doses which induce  
15 release of pituitary MIF into the circulation), and in animals challenged with TNF $\alpha$  (i.e., where the exogenous TNF $\alpha$  overrides the beneficial reduction in circulating endogenous TNF $\alpha$  concentrations achieved using anti-MIF). The ability of anti-MIF therapy to  
20 protect animals challenged with TNF $\alpha$  indicates that neutralization of MIF during the later, post-acute phase of septic shock is efficacious. The protective effect of the antibody may be attributed, in part, to neutralization of pituitary and macrophage MIF  
25 released in the post-acute phase of septic shock when circulating MIF is readily detected. Because MIF is a necessary component of the shock syndrome, and the peak of serum MIF expression occurs after that of TNF $\alpha$ , MIF inhibitors may be used to successfully treat  
30 cytokine-mediated conditions even later than the point at which administration of TNF $\alpha$  inhibitors is no longer effective.

Since steroids are potent inhibitors of cytokine production, their effects were examined on MIF  
35 secretion by macrophages and pituitary cells (see

Section 13, infra). Surprisingly, steroids were found to induce rather than inhibit MIF secretion by these cells. The secretion of MIF in response to steroids may reduce the benefit of steroid therapy currently  
5 used to treat inflammation. Therefore, MIF inhibition therapy may be used in conjunction with steroids to treat shock and other cytokine-mediated pathological states, particularly in chronic inflammatory states such as rheumatoid arthritis. Such combination  
10 therapy may be beneficial even subsequent to the onset of pathogenic or other inflammatory responses. For example, administration of steroids alone inhibits the TNF $\alpha$  response only if given simultaneously with or before LPS challenge. In the clinical setting, the  
15 administration of steroids subsequent to the onset of septic shock symptoms has proven of little benefit (Bone et al., 1987, N. Engl. J. Med. 317: 653-658; Spring et al., 1984, N. Engl. J. Med. 311: 1137-1141). Combination steroids/MIF inhibition therapy may be  
20 used to overcome this obstacle. In another example, when treating conditions such as rheumatoid arthritis, the administration of steroids alone can result in the induction of MIF, which may override the inhibitory effects of the administered steroid on the  
25 inflammatory response. In such cases, MIF inhibition therapy can be used in conjunction with steroid treatment. Therapy can be designed to inhibit MIF release and/or activity locally and/or systemically.

Finally, in yet another aspect of the invention,  
30 newly identified MIF-inhibitors are described. For example, the Applicant's have discovered that certain steroid derivatives commonly thought to be inactive or thought to block the anti-inflammatory effects of steroids ("anti-steroids") actually inhibit MIF  
35 release in response to subsequent steroid challenge.

Such "inert" or "anti-steroid" compounds may be used in conjunction with anti-inflammatory steroids to treat inflammatory diseases, especially non-infectious inflammatory diseases such as autoimmunity, rheumatoid arthritis, graft versus host disease etc. In particular, the data described herein (see Section 13, infra) show that steroids, such as dexamethasone, which are commonly used to treat inflammation actually induce MIF release. The steroid-induced release of MIF occurs at dose ranges that are used in vivo to avoid side effects, e.g., severe water retention, etc. (i.e., higher doses of steroids which result in serious side effects, and lower doses of steroids which are ineffective to treat inflammation do not appear to induce MIF release). The release of MIF in response to steroid challenge can be inhibited by pretreating with a MIF-inhibiting dose of the so-called "inactive" or "anti-steroid" derivatives, such as 20 $\alpha$ -dihydrocortisol. Because these derivatives are "inert" high doses which inhibit-MIF release should be well tolerated. The prior administration of the "inactive" compound should not induce side effects, should inhibit MIF release, and thus, may potentiate the anti-inflammatory action of the "active" steroid subsequently administered. Assays are described herein to identify additional compounds that can be used to inhibit the steroid-induced release of MIF from cellular sources in the body.

The following subsections summarize the data presented in Sections 6 through 15, infra.

#### 5.1.1. IDENTIFICATION OF MIF AS A PITUITARY HORMONE

To investigate the production and release of novel pituitary products, we examined the secretory profile of the murine anterior pituitary cell line,

AtT-20. These corticotrophic cells have been used widely to study ACTH release and we thought they were potentially useful cells in which to begin to investigate humoral interactions between the central nervous system (CNS), the hypothalamus and the periphery. As shown in Section 9, infra, we incubated AtT-20 cells with LPS for various time intervals and examined culture supernatants by SDS-PAGE. These analyses revealed the specific release by pituitary cells of an apparently novel 12.5 kDa protein. The 12.5 kDa protein was then isolated and determined by amino-terminal sequencing to be highly similar to an amino acid sequence predicted from a recently cloned human MIF cDNA (Weiser et al., 1989, supra). We cloned this pituitary protein from the cDNA of LPS-stimulated AtT-20 cells and showed that it is actually the murine homolog of human MIF (90% identity over 115 amino acids) (see Section 6, infra). As little as 100 pg/ml of LPS was found to induce pituitary cell secretion of MIF. However, pituitary MIF secretion was not induced by TNF- $\alpha$ , IL-1 $\beta$ , IL-6 or IFN- $\gamma$  (see Section 11, infra).

Immunocytochemical analyses revealed that resting, nonstimulated pituitary cells contain significant amounts of preformed MIF. Thus, a large fraction of the MIF that is released by pituitary cells in response to LPS arises from stored, intracellular pools. Electron microscopic studies of whole mouse pituitaries labeled with immunogold-conjugated anti-MIF antibody localize MIF to granules present within corticotrophic cells. It appears that in vivo, MIF-containing granules are released either by the direct action of circulating endotoxin or by specific hypothalamic releasing factor(s).

35

In vivo studies in mice showed that pituitary MIF mRNA levels increase after LPS challenge and reach a plateau after 16-24 hours. Over the same time course, the pituitary content of preformed MIF protein

5 decreases to almost undetectable levels. To determine whether the release of pituitary MIF is associated with a concomitant rise in serum MIF, we obtained blood samples from LPS-injected mice and analyzed serum for MIF content by western blotting. In normal

10 mice, MIF was detected 2 hours after administration of LPS and its levels increased in serum over approximately 20 hours. This time course for the rise in serum MIF is consistent both with the decrease in pituitary MIF protein and with the increase in

15 pituitary MIF mRNA that occurred after injection of LPS. In contrast, MIF in the serum of hypophysectomized mice showed a markedly different pattern of induction. Serum specimens obtained from LPS-treated, hypophysectomized mice showed no

20 detectable MIF at 20 hours, the time at which serum MIF levels were highest in normal mice. These data are consistent with the notion that the pituitary is critical to the MIF response that appears in serum during endotoxemia, and that pituicytes contribute to

25 serum levels of MIF. We then calculated the amount of preformed MIF stored in the pituitary and the amount of MIF that appears in serum after activation by LPS of the hypothalamic-pituitary axis. The quantities of MIF present in the pituitary and in serum were well

30 within the range observed for ACTH and other anterior pituitary hormones (Table I).



**TABLE I**

**COMPARISON OF PITUITARY MACROPHAGE MIGRATION  
INHIBITORY FACTOR (MIF) CONTENT AND SERUM MIF LEVELS  
WITH LEVELS OF OTHER ANTERIOR PITUITARY HORMONES**

5	Hormone	Pituitary content <sup>a</sup> (% of total protein)	Serum level <sup>b</sup> (ng ml <sup>-1</sup> )
	MIF	0.05	80-340
10	ACTH	0.2	2-500
	Prolactin	0.08	200-300

<sup>a</sup> Pituitary hormone content at resting conditions.

15 <sup>b</sup> Serum hormone level under conditions of maximal  
pituitary stimulation.

#### 5.1.2. MACROPHAGE MIF PRODUCTION

As shown in Section 12, infra, hypophysectomized  
20 mice have a markedly abnormal serum MIF response  
compared with control mice. The gradual increase in  
serum MIF over 20 hours is not observed. Instead, on  
closer examination, a peak of serum MIF is observed  
25 during the early, acute phase of endotoxaemia ( $\leq 3$  h  
after LPS injection). Given the lack of a pituitary  
in hypophysectomized mice, and the well-accepted  
unresponsiveness of T cells to LPS, we considered that  
the MIF present in the circulation at 2 hours must  
reflect the release of this cytokine by an additional  
30 LPS-sensitive cell population. Based on the  
observation that such an early peak in the serum MIF  
response is reminiscent of a macrophage TNF $\alpha$  response,  
we examined the possibility that MIF is also produced  
by cells of the monocyte-macrophage lineage.

35 Significant amounts of preformed MIF protein were  
found in the resting, nonstimulated murine monocyte

cell line RAW 264.7, murine peritoneal macrophages and the human monocyte cell line THP-1. Western blotting revealed that the MIF content of murine or human monocytes-macrophages was similar to that of the murine ASL-1 and the human Jurkat T cell lines. In contrast, cell lysates obtained from purified polymorphonuclear leukocytes did not contain detectable MIF protein. Whole tissue analyses demonstrated significant amounts of both MIF protein and mRNA in organs that have a high content of macrophages (see Section 14, *infra*), complementing recent observations that MIF mRNA is present constitutively in tissues such as the spleen, the liver, and the kidney. (Lanahan et al., 1992, Mol. Cell. Biol. 12: 3919-3929).

Various proinflammatory stimuli, such as LPS, TNF $\alpha$  and IFN- $\gamma$  were observed to be potent inducers of macrophage MIF release (see Section 12, *infra*). Secretion of significant amounts of MIF occurred at LPS concentrations (10-100 pg ml<sup>-1</sup>) that are lower in general than has been observed to be optimal for the induction of TNF $\alpha$  release. Significant amounts of preformed MIF mRNA were also present in resting macrophages, and stimulation by LPS increased these mRNA levels approximately two-fold. Peak mRNA stimulation also occurred at very low concentrations of LPS (1 pg ml<sup>-1</sup>).

#### 5.1.3. CLONING, EXPRESSION AND BIOCHEMICAL CHARACTERIZATION OF MIF

The cloning of murine MIF (muMIF) from an anterior pituitary cell line and the purification of native MIF from mouse liver are described in Section 6, *infra*. For comparison purposes, human MIF (huMIF) was cloned from the Jurkat T-cell line. Sequence analysis of murine pituitary MIF and human T-cell MIF

cdNA demonstrated a 90% homology between the murine and human proteins. Cloned Jurkat T-cell MIF cdNA differs by one nucleotide from the first published human T-cell MIF cdNA (Weiser et al., 1989), but is in  
5 agreement with human MIF cdNA derived from human lens tissue (Wistow et al., 1993, Proc. Natl. Acad. Sci. USA 90: 1272-1275), and with a T-cell glycosylation inhibition factor (GIF) cdNA (Mikayama et al., 1993, Proc. Natl. Acad. Sci. USA 90: 10056-10060). The  
10 single nucleotide difference results in a change in the deduced amino acid sequence (Ser 106 → Asn 106), thus increasing homology of the human and murine proteins.

DNA homology analysis showed that both human and  
15 murine MIF lack a conventional N-terminal leader sequence. Thus, MIF joins a growing list of cytokines, such as IL-1 (Rubartelli et al., 1990, EMBO J. 9: 1503-1510), basic fibroblast growth factor (bFGF; Jackson et al., 1992, Proc. Natl. Acad. Sci.  
20 USA 89: 10691-10695), and a secreted form of cyclophilin (Sherry et al., 1992, Proc. Natl. Acad. Sci. USA 89: 3511-3515) which are released from cells by non-classical protein secretion pathways.

Recombinant murine and human MIF were expressed  
25 in *E. coli*, purified to homogeneity by a simple two-step procedure, and the proteins characterized by a variety of biochemical and biological criteria. Native MIF obtained from mouse liver was found to be monomeric and to be free of significant post-  
30 translational modifications as assessed both by endoglycosidase F treatment followed by SDS-PAGE/Western blotting and by mass spectroscopic analysis. Recombinant and native MIF exhibited a number of comparable effects on monocytes/macrophages.  
35 Recombinant MIF was found to inhibit monocyte

migration when assayed in modified Boyden chambers. MIF also induced TNF $\alpha$  secretion, and promoted the release of nitric oxide (NO) from macrophages primed with IFN- $\gamma$ . Circular dichroism (CD) spectroscopy  
5 revealed that bioactive, recombinant MIF exhibits a highly ordered three-dimensional structure, with a > 55% content of  $\beta$ -pleated sheet and  $\alpha$ -helix. Thermodynamic stability studies also showed that despite a high content of ordered secondary structure  
10 elements, the conformation of MIF was readily perturbed by denaturing solvent conditions. These studies define a number of the biochemical and biological properties of recombinant and purified, native MIF and provide, for the first time,  
15 information concerning the three-dimensional structure of this protein.

#### 5.1.4. BIODISTRIBUTION OF MIF AND MIF RECEPTORS

The working example in Section 14, infra,  
20 presents a study that illustrates, the distribution of MIF and MIF-binding sites, i.e. MIF-receptors within the body. The study indicates that MIF receptors are present in kidney cells, for example. Further, two such MIF receptors are identified, for the first time,  
25 in the working example presented, infra, in Section 13. Such receptors may be genetically engineered for use as MIF inhibitors.

#### 5.2. INHIBITORS OF MIF ACTIVITY

30 Described below are factors which may be used as MIF antagonists, i.e., factors which inhibit the biologic activity of MIF.

35

### 5.2.1. MIF-BINDING PARTNERS

Factors that bind MIF and neutralize its biological activity, hereinafter referred to as MIF binding partners, may be used in accordance with the invention as treatments of conditions involving cytokine-mediated toxicity. While levels of MIF protein may increase due to endotoxin challenge, the interaction of inhibitory MIF-binding partners with MIF protein prohibits a concomitant increase in MIF activity. Such factors may include, but are not limited to anti-MIF antibodies, antibody fragments, MIF receptors, and MIF receptor fragments.

Various procedures known in the art may be used for the production of antibodies to epitopes of recombinantly produced (e.g., using recombinant DNA techniques described infra), or naturally purified MIF. Neutralizing antibodies, i.e. those which compete for or sterically obstruct the binding sites of the MIF receptor are especially preferred for diagnostics and therapeutics. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by an Fab expression library.

For the production of antibodies, various host animals may be immunized by injection with MIF and/or a portion of MIF. Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants

such as BCG (bacille Calmette-Guerin) and Corynebacterium parvum.

Monoclonal antibodies to MIF may be prepared by using any technique which provides for the production  
5 of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Kohler and Milstein, (Nature, 1975, 256:495-497), the human B-cell hybridoma technique (Kosbor et al., 1983,  
10 Immunology Today, 4:72; Cote et al., 1983, Proc. Natl. Acad. Sci., 80:2026-2030) and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). In addition, techniques developed for the production  
15 of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen  
20 specificity together with genes from a human antibody molecule of appropriate biological activity can be used. Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce MIF-specific  
25 single chain antibodies.

Antibody fragments which recognize specific MIF epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')<sub>2</sub> fragments which can be produced by  
30 pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-  
35 1281) to allow rapid and easy identification of

monoclonal Fab fragments with the desired specificity to MIF.

MIF receptors, MIF receptor fragments, and/or MIF receptor analogs may, in accordance with the invention, be used as inhibitors of MIF biological activity. By binding to MIF protein, these classes of molecules may inhibit the binding of MIF to cellular MIF receptors, thus disrupting the mechanism by which MIF exerts its biological activity. Small organic molecules which mimic the activity of such molecules are also within the scope of the present invention.

MIF receptors may include any cell surface molecule that binds MIF in an amino acid sequence-specific and/or structurally-specific fashion. Such MIF receptors include, but are not limited to the 72 kD MIF receptor class having a partial amino acid sequence of:

AKKGAVGGI

and the 52 kD receptor having a partial amino acid sequence of:

I-X-HNTVATEI(S)(G)YN(N/G)A(M)

both of which are presented in the working example in Section 13, infra. The residues in parenthesis are provisional assignment.

Additional MIF receptors and genes that encode MIF receptors may be identified, isolated, and cloned using a variety of techniques well known to those of ordinary skill in the art. For example, MIF receptor molecules may be identified and isolated using standard affinity chromatography techniques wherein those molecules exhibiting sequence- and/or structural binding specificity to MIF protein are separated from other non-MIF binding molecules. The MIF binding proteins may be additionally purified, using standard

35

techniques, at which point the protein may be tested and utilized for its ability to inhibit MIF.

Alternatively, the amino acid sequence of the purified protein may be at least partially determined, and then used to design oligonucleotides with which to screen cDNA and/or genomic libraries in order to clone the gene(s) that encode(s) MIF receptors, techniques of which are well known to those of skill in the art. Further, new MIF receptor genes may be cloned by construction of a cDNA library in a mammalian expression vector such as pcDNA1, that contains SV40 origin of replication sequences which permit high copy number expression of plasmids when transferred into COS cells. The expression of the MIF receptor on the surface of transfected COS cells may be detected in a number of ways, including the use of radioactive, fluorescent, or enzymatically labeled MIF. Cells expressing an MIF receptor may be enriched by subjecting transfected cells to a FACS (fluorescent activated cell sorter). For a review of cloning strategies which may be used, see e.g., Maniatis, 1989, Molecular Cloning, A Laboratory Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, (Green Publishing Associates and Wiley Interscience, N.Y.)

Fragments of any of the MIF receptors described above may also be used as MIF inhibitory agents, and any MIF receptor fragment possessing any amino, carboxy, and/or internal deletion that specifically binds MIF so as to inhibit MIF biological activity is intended to be within the scope of this invention. An amino and/or carboxy deletion refers to a molecule possessing amino and/or carboxy terminal truncations of at least one amino acid residue. An internal deletion refers to molecules that possess one or more



non-terminal deletions of at least one amino acid residue. Among these MIF receptor fragments are truncated receptors in which the cytoplasmic or a portion of the cytoplasmic domain has been deleted,  
5 and fragments in which the cytoplasmic and the transmembrane domain(s) has been deleted to yield a soluble MIF receptor containing all or part of the MIF receptor extracellular domain.

MIF receptor analogs which specifically bind MIF  
10 may also be used to inhibit MIF activity. Such MIF receptor analogs may include MIF receptor or receptor fragments further possessing one or more additional amino acids located at the amino terminus, carboxy terminus, or between any two adjacent MIF receptor  
15 amino acid residues. The additional amino acids may be part of a heterologous peptide functionally attached to all or a portion of the MIF receptor protein to form a MIF receptor fusion protein. For example, and not by way of limitation, the MIF  
20 receptor, or a truncated portion thereof, can be engineered as a fusion protein with a desired Fc portion of an immunoglobulin. MIF receptor analogs may also include MIF receptor or MIF receptor fragments further possessing one or more amino acid  
25 substitutions of a conservative or non-conservative nature. Conservative amino acid substitutions consist of replacing one or more amino acids with amino acids of similar charge, size, and/or hydrophobicity characteristics, such as, for example, a glutamic acid  
30 (E) to aspartic acid (D) amino acid substitution. Non-conservative substitutions consist of replacing one or more amino acids with amino acids possessing dissimilar charge, size, and/or hydrophobicity characteristics, such as, for example, a glutamic acid  
35 (E) to valine (V) substitution.

The MIF receptors, MIF receptor fragments and/or analogs may be made using recombinant DNA techniques. Here, the nucleotide sequences encoding the peptides of the invention may be synthesized, and/or cloned,  
5 and expressed according to techniques well known to those of ordinary skill in the art. See, for example, Ausubel, F.M. et al., eds., 1989, Current Protocols In Molecular Biology, Vol. 1 and 2, John Wiley and Sons, New York. Caruthers, et al., 1980, Nuc. Acids Res.  
10 Symp. Ser. 7:215-233; Crea and Horn, 1980, Nuc. Acids Res. 9(10):2331; Matteucci and Caruthers, 1980, Tetrahedron Letters 21:719; and Chow and Kempe, 1981, Nuc. Acids Res. 9(12):2807-2817.

Alternatively, the protein itself could be  
15 produced using chemical methods to synthesize the amino acid sequence in whole or in part. For example, peptides can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography. (E.g., see  
20 Creighton, 1983, Proteins Structures And Molecular Principles, W.H. Freeman and Co., N.Y. pp. 50-60.) The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, 1983,  
25 Proteins, Structures and Molecular Principles, W.H. Freeman and Co., N.Y., pp. 34-49).

These molecules may also be synthesized utilizing alternative procedures which may advantageously affect certain of the molecules' properties, such as  
30 stability, bioavailability, and MIF inhibitory activity. For example, MIF receptors, MIF receptor fragments, and MIF receptor analogs may be synthesized such that one or more of the bonds which link the amino acid residues of the peptides are non-peptide  
35 bonds. These alternative non-peptide bonds may be

formed by utilizing reactions well known to those in the art, and may include, but are not limited to imino, ester, hydrazide, semicarbazide, and azo bonds, to name but a few. In another embodiment, the

5 proteins may be synthesized with additional chemical groups present at their amino and/or carboxy termini. For example, hydrophobic groups such as carbobenzoxyl, dansyl, or t-butyloxycarbonyl groups, may be added to the peptides' amino terminus. Further, the peptides

10 may be synthesized such that their steric configuration is altered. For example, the D-isomer of one or more of the amino acid residues of the peptide may be used, rather than the usual L-isomer.

15 5.2.2. MIF-RECEPTOR ANTAGONISTS

Molecules which inhibit MIF biological activity by binding to MIF receptors may also be utilized for the treatment of conditions involving cytokine-mediated toxicity. Such molecules may include, but

20 are not limited to anti-MIF receptor antibodies and MIF analogs.

Anti-MIF receptor antibodies may be raised and used to neutralize MIF receptor function. Antibodies against all or any portion of a MIF receptor protein

25 may be produced, for example, according to the techniques described in Section 5.2.1., supra.

MIF analogs may include molecules that bind the MIF receptor but do not exhibit biological activity. Such analogs compete with MIF for binding to the MIF

30 receptor, and, therefore, when used in vivo, may act to block the effects of MIF in the progress of cytokine-mediated toxicity. A variety of techniques well known to those of skill in the art may be used to design MIF analogs. The coding sequence for human MIF

35 described herein, differs from the published sequence,

resulting in a serine to asparagine change in the gene product at amino acid residue number 106. Both the corrected human sequence and that of the murine protein is described in Section 6, infra, and shown in FIG. 2. Recombinant DNA techniques may be used to produce modified MIF proteins containing, for example, amino acid insertions, deletions and/or substitutions which yield MIF analogs with receptor binding capabilities, but no biological activity.

Alternatively, MIF analogs may be synthesized using chemical methods such as those described above in Section 5.2.1. (see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, N.Y.).

MIF receptors and/or cell lines that express MIF receptors may be used to identify and/or assay potential MIF antagonists. For example, one method that may be pursued in the identification of such MIF antagonist molecules would comprise attaching MIF receptor molecules to a solid matrix, such as agarose or plastic beads, microtiter wells, or petri dishes, using techniques well known to those of skill in the art, and subsequently incubating the attached MIF receptor molecules in the presence of a potential MIF analog or analogs. After incubating, unbound compounds are washed away, and the MIF receptor-bound compounds are recovered. In this procedure, large numbers of types of molecules may be simultaneously screened for MIF receptor-binding activity. Bound molecules may be eluted from the MIF receptor molecules by, for example, competing them away from the MIF receptor molecules with the addition of excess MIF, changing the pH or osmotic strength of the buffer or adding chaotropic agents. The binding/elution

35

steps bring about a partial purification of the molecules of interest.

In order to continue the purification process, the eluted molecules may be further fractionated by one or more chromatographic and/or other separation techniques well known in the art until the molecule(s) of interest is (are) purified to the extent necessary. Once compounds having MIF-receptor binding capabilities are identified, the compounds may be assayed for their effects on cytokine-mediated toxicity using cell lines such as those described in this Section, or by normal experimental animal models or alternatively, by utilizing transgenic animal models such as those described in Section 5.4, infra.

Alternatively, screening of peptide libraries with recombinantly produced MIF receptors and/or MIF receptor fragments may be used to identify potential MIF analogs. Once peptides that bind MIF receptor are identified using this screening technique, their effects on cytokine-mediated toxicity may be assayed using cells lines such as those described in this Section, or alternatively, may be evaluated using normal experimental animal models or transgenic animals such as those described in Section 5.4., infra. Small organic molecules which mimic the activity of such peptides are also within the scope of the present invention.

Random peptide libraries consist of all possible combinations of amino acids, and such libraries may be attached to a suitable small particulate solid phase support and used to identify peptides that are able to bind to a given receptor (Lam, K.S. et al., 1991, Nature 354: 82-84). The screening of peptide libraries may have therapeutic value in the discovery of pharmaceutical agents that act to inhibit the

biological activity of receptors through their interactions with the given receptor.

Identification of molecules that are able to bind to the MIF receptor may be accomplished by screening a peptide library with recombinant soluble MIF receptor protein. Methods for expression and purification of molecules such as MIF receptors are well known to those of skill in the art. For screening, it is preferred to label or "tag" the MIF receptor molecule. The protein may be conjugated to enzymes such as alkaline phosphatase or horseradish peroxidase or to other reagents such as fluorescent labels which may include fluorescein isothiocyanate (FITC), phycoerythrin (PE) or rhodamine. Conjugation of any given label to the MIF receptor may be performed using techniques that are routine in the art. Alternatively, MIF receptor expression vectors may be engineered to express a chimeric MIF receptor protein containing an epitope for which a commercially available antibody exists. The epitope-specific antibody may be tagged using methods well known in the art including labeling with enzymes, fluorescent dyes or colored or magnetic beads.

The "tagged" MIF receptor or receptor/conjugate is incubated with the random peptide library for 30 minutes to one hour at 22°C to allow complex formation between MIF receptor and peptide species within the library. The library is then washed to remove any unbound MIF receptor protein. If MIF receptor has been conjugated to alkaline phosphatase or horseradish peroxidase the whole library is poured into a petri dish containing a substrates for either alkaline phosphatase or peroxidase, for example, 5-bromo-4-chloro-3-indoyl phosphate (BCIP) or 3,3',4,4'-diaminobenzidine (DAB), respectively. After

incubating for several minutes, the peptide/solid phase-MIF receptor complex changes color, and can be easily identified and isolated physically under a dissecting microscope with a micromanipulator. If a fluorescent tagged MIF receptor molecule has been used, complexes may be isolated by fluorescence-activated sorting. If a chimeric MIF protein expressing a heterologous epitope has been used, detection of the peptide/MIF receptor complex may be accomplished by using a labeled epitope-specific antibody. Once isolated, the MIF receptor conjugate may be eluted off, the peptide support washed, and the identity of the peptide attached to the solid phase support determined by peptide sequencing.

MIF analogs may also be identified using cell lines that express MIF receptor. Such cell lines may be ones which naturally express the receptor, such as RAW 264.7 cells, or alternatively, cell lines that have been engineered using recombinant techniques to express MIF receptor. These cell lines may also be used to evaluate potential MIF analogs identified using MIF receptor binding techniques such as those described above.

With respect to engineered cell lines, a variety of cells may be utilized as host cells for expression of the recombinant MIF receptor, including, but not limited to animal cell systems infected with recombinant virus expression vectors (e.g., adenovirus, vaccinia virus) including cell lines engineered to contain multiple copies of the MIF receptor DNA either stably amplified (e.g., CHO/dhfr) or unstably amplified in double-minute chromosomes (e.g., murine cell lines). In cases where an adenovirus is used as an expression vector, the MIF receptor coding sequence may be ligated to an adenovirus transcription-

translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a

5 non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing MIF receptor in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. (USA) 81:3655-3659). Alternatively,

10 the vaccinia 7.5K promoter may be used. (See, e.g., Mackett et al., 1982, Proc. Natl. Acad. Sci. (USA) 79: 7415-7419; Mackett et al., 1984, J. Virol. 49: 857-864; Panicali et al., 1982, Proc. Natl. Acad. Sci. 79: 4927-4931).

15 Specific initiation signals may also be required for efficient translation of inserted MIF receptor coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where the entire MIF receptor gene, including its own

20 initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional transcriptional control signals may be needed. However, in cases where only a portion of the MIF receptor coding sequence is inserted, exogenous

25 transcriptional control signals, including the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the MIF receptor coding sequence to ensure translation of the entire insert. These exogenous

30 translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc.

35



(see Bittner et al., 1987, Methods in Enzymol. 153: 516-544).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cells lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, and for any normal glycosylation, and/or phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, WI38, etc.

The cell lines may be utilized to screen and identify MIF analogs. Synthetic compounds, natural products, and other sources of potentially biologically active materials can be screened in a number of ways, by, for example, testing a compound's ability to inhibit binding of MIF to a MIF receptor. Standard receptor binding techniques may be utilized for this purpose.

The ability of anti-MIF receptor antibodies and potential MIF analogs to reduce or inhibit MIF biological activity may be assayed in vivo by utilizing animals expressing MIF receptor, for instance, normal animals. Such animals may also include transgenic animal models such as those described below in Section 5.4, infra.

### 5.2.3. OTHER INHIBITORS OF MIF ACTIVITY

As explained in Section 5.1, supra, certain steroids, commonly thought to be either inactive or "anti-steroidal" actually inhibit the release of MIF;   
5 e.g., 20 $\alpha$  dihydrocortisol. These steroids, or any other compound which inhibits the release of preformed MIF, can be used in combination therapy with anti-inflammatory steroids.

Compounds which inhibit the release of MIF can be   
10 identified in cell based assays, such as the one described in Section 13, infra. In general, any pituitary or macrophage cell line that releases MIF in response to a challenge dose of steroid can be used. The assay can be conducted by adding the test compound   
15 to the cells in culture which are then challenged with a dose of steroid known to induce MIF release. Test compounds may be administered simultaneously with, or up to several hours before or after the challenge dose so as to identify agents that are useful in inhibiting   
20 the MIF response at different stages, i.e., inhibiting release of pre-formed MIF, versus inhibiting de novo synthesis and release, versus inhibiting both.

The conditioned media is then collected from the cultured cells and assayed for MIF; e.g., by   
25 immunoassay, including but not limited to an ELISA, Western blot, radioimmunoassay, etc. A reduced amount of MIF in the conditioned media indicates that the test compound inhibits the steroid-induced release of MIF. Compounds so identified in this assay may be   
30 used in combination therapy with steroids to treat inflammation. "Biologically inert" or innocuous compounds, such as the inactive steroids, or steroids which can be used at doses that do not cause undesired side effects, may be preferred for therapeutic use.   
35 However, any inhibitory compounds having a good

therapeutic index, e.g., low toxicity and little or no side effects may be used.

#### 5.2.4. DOSE AND TREATMENT REGIMENS

- 5       Inhibitors of MIF biological activity such as anti-MIF antibodies, MIF receptors, MIF receptor fragments, MIF receptor analogs, anti-MIF receptor antibodies, MIF analogs and inhibitors of MIF release, may be administered using techniques well known to
- 10   those in the art. Preferably, agents are formulated and administered systemically. Techniques for formulation and administration may be found in "Remington's Pharmaceutical Sciences", 18th ed., 1990, Mack Publishing Co., Easton, PA. Suitable routes may
- 15   include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal,
- 20   intranasal, or intraocular injections, just to name a few. Most preferably, administration is intravenous. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks'
- 25   solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.
- 30       Effective concentrations and frequencies of dosages of the MIF inhibitory compounds invention to be administered may be determined through procedures well known to those in the art, which address such parameters as biological half-life, bioavailability,
- 35   and toxicity. In the case of anti-MIF antibodies, a

preferred dosage concentration may range from about 0.1 mg/kg body weight to about 20 mg/kg body weight, with about 10 mg/kg body weight being most preferred. Because antibodies typically exhibit long half-lives  
5 in circulation, a single administration of antiserum may be sufficient to maintain the required circulating concentration. In the case of compounds exhibiting shorter half-lives, multiple doses may be necessary to establish and maintain the requisite concentration in  
10 circulation.

MIF inhibitors may be administered to patients alone or in combination with other therapies. Such therapies include the sequential or concurrent administration of inhibitors or antagonists of  
15 initiators of cytokine-mediated toxicity (e.g. anti-LPS), inhibitors or antagonists of participants in the endogenous cytokine responses (e.g. anti-TNF $\alpha$ , anti-IL-1, anti-IFN- $\gamma$ , IL-1 RA); and compounds that inhibit or antagonize cytokine-mediated toxicity directly  
20 (e.g. steroids, glucocorticoids or IL-10). MIF inhibitor dosage concentration and frequency may be altered when used as a part of a combination therapy and, therefore appropriate tests must be performed in order to determine the best dosage when more than one  
25 class of inhibitory compounds is to be administered.

Because MIF expression reaches a peak level in response to endotoxin challenge later than TNF $\alpha$ , anti-MIF compounds may be administered after the period within which anti-TNF $\alpha$  inhibitors are effective. As  
30 shown in the working example, described in Section 7, infra, anti-MIF antibody conferred full protection against endotoxemia in animals. Anti-MIF has also been shown in Applicants' pilot studies to reduce circulating TNF $\alpha$  levels, indicating that anti-MIF  
35 inhibits the proinflammatory spectrum of activity of

MIF, and that anti-MIF inhibits the inflammatory cytokine cascade generally. In in vivo experiments, moreover, anti-MIF was shown to protect against lethal shock induced by administration of exogenous TNF $\alpha$  (see  
5 Section 7, infra). Therefore, the beneficial effect of the anti-MIF antibody probably resides in its ability to neutralize the bioactivity of both macrophage MIF released during the acute phase in response to a proinflammatory stimulus (e.g., LPS, or  
10 TNF $\alpha$  and IFN- $\gamma$ ), and MIF released by both the pituitary and macrophages during the post-acute phase of the shock response.

15 5.3. INHIBITORS OF MIF AND/OR  
MIF RECEPTOR GENE EXPRESSION

Nucleotide sequences derived from the coding, non-coding, and/or regulatory sequences of the MIF and/or MIF receptor genes may be used to prevent or reduce the expression of these genes, leading to a  
20 reduction or inhibition of MIF activity. The nucleotide sequence encoding the human MIF protein has been reported. In addition, as presented in the working example in Section 6, infra, the nucleotide sequence of human MIF has been corrected, and a cDNA  
25 corresponding to the nucleotide sequence encoding the murine MIF protein has now been identified. Further, the MIF receptor amino acid sequence provided in the working example in Section 15, infra, may, for example, be used to design oligonucleotides for the  
30 regulation of MIF receptor genes. Among the techniques by which such regulation of gene expression may be accomplished are, as described below, antisense, triple helix, and ribozyme approaches. Administration of these nucleotide sequences,  
35 therefore, may be used to temporarily block expression and/or transcription of the MIF and/or MIF receptor

genes as one method of treatment for conditions involving cytokine-mediated toxicity.

These approaches which target gene expression may be used alone, in combination with each other, or  
5 alternatively, in conjunction with any of the inhibitory MIF-binding and/or MIF receptor antagonist procedures described above. Further, these gene regulation approaches may be used together with anti-TNF $\alpha$ , anti-initiators and/or other anti-cytokine  
10 therapies.

#### 5.3.1. ANTI-SENSE RNA AND RIBOZYMES

Within the scope of the invention are oligo-ribonucleotide sequences, that include anti-sense RNA  
15 and DNA molecules and ribozymes that function to inhibit the translation of MIF and/or MIF receptor mRNA. Anti-sense RNA and DNA molecules act to directly block the translation of mRNA by binding to targeted mRNA and preventing protein translation,  
20 either by inhibition of ribosome binding and/or translocation or by bringing about the nuclease degradation of the mRNA molecule itself.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific  
25 hybridization of the ribozyme molecule to complementary target RNA, followed by a endonucleolytic cleavage. Within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically  
30 and efficiently catalyze endonucleolytic cleavage of MIF and/or MIF receptor mRNA sequences.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage  
35 sites which include the following sequences, GUA, GUU

and GUC (See FIG. 2 for an illustration of such potential sites on murine and human MIF cDNA). Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for predicted structural features such as secondary structure that may render the oligonucleotide sequence unsuitable. The suitability of candidate targets may also be evaluated by testing their accessibility to hybridization with complimentary oligonucleotides, using ribonuclease protection assays.

Both anti-sense RNA and DNA molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides well known in the art such as, for example, solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

Various modifications to the DNA molecules may be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not limited to the addition of flanking sequences of ribo- or deoxy- nucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phospho-

diesterase linkages within the oligodeoxyribonucleotide backbone.

#### 5.3.2. TRIPLEX DNA FORMATION

- 5 Oligodeoxyribonucleotides can form sequence-specific triple helices by hydrogen bonding to specific complementary sequences in duplexed DNA. Interest in triple helices has focused on the potential biological and therapeutic applications of
- 10 these structures. Formation of specific triple helices may selectively inhibit the replication and/or gene expression of targeted genes by prohibiting the specific binding of functional trans-acting factors.
- Oligonucleotides to be used in triplex helix
- 15 formation should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either
- 20 purines or pyrimidines to be present on one strand of a duplex. Oligonucleotide sequences may be pyrimidine-based, which will result in TAT and CGC triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich
- 25 oligonucleotides provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, oligonucleotides may be chosen that are purine-rich, for example, containing a stretch of G residues.
- 30 These oligonucleotides will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.
- 35 Alternatively, the potential sequences that can be



targeted for triple helix formation may be increased by creating a so called "switchback" oligonucleotide. Switchback oligonucleotides are synthesized in an alternating 5'-3', 3'-5' manner, such that they base  
5 pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

10           5.3.3. ADMINISTRATION OF OLIGONUCLEOTIDES

For anti-MIF therapeutic uses, the inhibitory oligonucleotides may be formulated and administered through a variety of means, including systemic, and localized, or topical, administration. Techniques for  
15 formulation and administration may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. The mode of administration may be selected to maximize delivery to a desired target organ in the body. For example, <sup>125</sup>I-  
20 MIF binding studies detailed in Section 12, infra, indicate that MIF is preferentially distributed to the liver and kidney. Therefore, oligonucleotides designed to inhibit expression of the MIF-receptor may be formulated for targeting to these organs; in this  
25 regard, liposome-encapsulated oligonucleotides may prove beneficial. Alternatively, MIF itself is produced in T cells and macrophages and, in response to endotoxin induction, in the pituitary. Therefore, oligonucleotides designed to inhibit the expression of  
30 MIF should be formulated for targeting to these organs.

For systemic administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous. For injection, the  
35 oligonucleotides of interest are formulated in aqueous

solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. In addition, the oligonucleotides may be formulated in solid or  
5 lyophilized form, then redissolved or suspended immediately prior to use. Systemic administration may also be accomplished by transmucosal, transdermal, or oral means. For transmucosal or transdermal administration, penetrants appropriate to the barrier  
10 to be permeated are used in the formulation. Such penetrants are generally known in the art. Transmucosal administration may be through nasal sprays or suppositories. For oral administration, oligonucleotides may be formulated into capsules,  
15 tablets, and tonics. For topical administration, the oligonucleotides of the invention are formulated into ointments, salves, gels, or creams, as is generally known in the art.

Alternatively, the oligonucleotides of the  
20 invention may first be encapsulated into liposomes, then administered as described above. Liposomes are spherical lipid bilayers with aqueous interiors. All molecules that are present in an aqueous solution at the time of liposome formation (in this case,  
25 oligonucleotides) are incorporated into this aqueous interior. The liposomal contents are both protected from the external microenvironment and, because liposomes fuse with cell membranes, are efficiently delivered into the cell cytoplasm, obviating the need  
30 to neutralize the oligonucleotides' negative charge.

The introduction of oligonucleotides into organisms and cells for such purposes may be accomplished by several means. For mammalian administration, each of the techniques described above  
35 for therapeutic oligonucleotide purposes may be used.

In addition, other standard techniques for introduction of nucleic acids into cells, including, but not limited to, electroporation, microinjection, and calcium phosphate precipitation techniques may be  
5 utilized.

#### 5.4. TRANSGENIC ANIMAL MODELS

Transgenic animals may be engineered, using techniques well known to those of skill in the art,  
10 whose cells may contain modified and/or additional MIF and or MIF receptor genes within their genomes. For example, animals may be produced that contain inactive MIF and/or MIF receptor genes, or alternatively, may contain additional MIF and/or MIF receptor genes.

15 Such transgenic animals may be used as model systems for the evaluation of cytokine responses in vivo, and may additionally serve as a means by which new drugs for the treatment of conditions involving cytokine-mediated toxicity are identified and tested.

20

##### 5.4.1. MIF AND/OR MIF RECEPTOR TRANSGENES

DNA containing the nucleotide coding sequence for all or any portion of the MIF gene may be used to produce transgenic animals. Alternatively, all or any  
25 portion of a gene encoding a MIF receptor may be used. Further, insertions, substitutions, and/or deletions of one or more nucleotides of the MIF and/or MIF receptor genes may also be utilized in the construction of the transgenic animals. Due to the  
30 degeneracy of the genetic code, other DNA sequences which encode substantially the same MIF or MIF receptor protein or a functional equivalent can also be used. The nucleotide coding sequence used to produce the transgenic animals of the invention may be  
35 regulated by any known promoter regulatory nucleotide

sequence. If it is required that expression of the transgene be limited to one or more specific tissues, tissue-specific enhancer regulatory sequences may also be used. Multiple copies of the genes or gene  
5 constructs may be stably integrated into the transgenic founder animals.

Any of the nucleotide coding sequences and/or regulatory sequences that will yield any of the variants described here can be produced using  
10 recombinant DNA and cloning methods which are well known to those of skill in the art. In order to produce the gene constructs used in the invention, recombinant DNA and cloning methods which are well known to those skilled in the art may be utilized (see  
15 Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, NY). In this regard, appropriate MIF and/or MIF receptor coding sequences may be generated from MIF cDNA or genomic clones using restriction enzyme sites  
20 that are conveniently located at the relevant positions within the sequences. Alternatively, or in conjunction with the method above, site directed mutagenesis techniques involving, for example, either the use of vectors such as M13 or phagemids, which are  
25 capable of producing single stranded circular DNA molecules, in conjunction with synthetic oligonucleotides and specific strains of *Escherichia coli* (*E. coli*) (Kunkel, T.A. et al., 1987, Meth. Enzymol. 154:367-382) or the use of synthetic  
30 oligonucleotides and PCR (polymerase chain reaction) (Ho et al., 1989, Gene 77:51-59; Kamman, M. et al., 1989, Nucl. Acids Res. 17: 5404) may be utilized to generate the necessary nucleotide coding sequences. Appropriate MIF and/or MIF receptor sequences may then  
35 be isolated, cloned, and used directly to produce

transgenic animals. The sequences may also be used to engineer the chimeric gene constructs that utilize regulatory sequences other than the MIF and/or MIF receptor promoter, again using the techniques

5 described here. These chimeric gene constructs would then also be used in the production of transgenic animals.

Transgenic animals may also be produced in which the function of the endogenous MIF and/or MIF receptor  
10 genes has been disrupted. To accomplish these endogenous gene disruptions, the technique of site-directed inactivation via gene targeting (Thomas, K.R. and Capecchi, M.R., 1987, Cell 51:503-512) may be used. Briefly, vectors containing some nucleotide  
15 sequences homologous to the endogenous gene of interest are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of, the nucleotide sequence of said endogenous gene.

20

#### 5.4.2. PRODUCTION OF TRANSGENIC ANIMALS

Animals of any species, including but not limited to mice, rats, rabbits, guinea pigs, pigs, mini-pigs, and non-human primates, e.g., baboons, squirrel  
25 monkeys and chimpanzees may be used to generate the transgenic animals of the invention. Any technique known in the art may be used to introduce the transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are  
30 not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No. 4,873,191); retrovirus-mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells  
35 (Thompson et al., 1989, Cell 56:313-321);

electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. (For a review of such techniques, see Gordon, 1989, 5 Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.)

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce 10 colonies of the particular genotype desired. Examples of such breeding strategies include but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order 15 to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic mice to produce mice homozygous for a given integration site in order to 20 both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; breeding animals to different inbred genetic backgrounds so as to examine 25 effects of modifying alleles on expression of the transgene.

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, 30 but not all their cells, i.e., mosaic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems.

35

### 5.5. DIAGNOSTIC APPLICATIONS

MIF protein and/or mRNA levels may be monitored in an individual, using standard techniques, as an indication of the deleterious aspects of a disease condition. For the measurement of MIF protein concentrations, such monitoring techniques include, but are not limited to immunological assays such as, for example, Western blots, immunoassays such as radioimmuno-precipitation, enzyme-linked immunoassays, and the like. For the measurement of MIF mRNA concentrations, such techniques may include, for example, hybridization techniques such as Northern blot analysis, or any RNA amplification techniques, which may involve, for example, polymerase chain reaction (the experimental embodiment set forth in Mullis, K.B., 1987, U.S. Patent No. 4,683,202) ligase chain reaction (Barany, F., 1991, Proc. Natl. Acad. Sci. USA 88: 189-193) self-sustained sequence replication (Guatelli, J.C. et al., 1990, Proc. Natl. Acad. Sci. USA 87: 1874-1878), transcriptional amplification system (Kwoh, D.Y. et al., 1989, Proc. Natl. Acad. Sci. USA 86: 173-1177), or Q-Beta Replicase (Lizardi, P.M. et al., 1988, Bio/Technology 6: 1197).

25

### 6. EXAMPLE: MOLECULAR CLONING OF MIF AND STRUCTURAL COMPARISON BETWEEN MURINE AND HUMAN MIF

30

#### 6.1. MATERIALS AND METHODS

##### 6.1.1. MATERIALS

Reagents for polymerase chain reaction (PCR), reverse transcription (RT), and other molecular biology techniques were purchased from Gibco BRL (Grand Island, NY) unless stated otherwise. PCR

35

buffer was from Perkin Elmer Cetus (Norwalk, CT); RNase inhibitor "rRNasin" was obtained from Promega (Madison, WI); RNAzol<sup>TM</sup>B was from TEL-TEST, INC. (Friendswood, TX); and oligonucleotides were purchased from OLIGOS ETC., INC. (Wilsonville, OR). Manual DNA sequencing was performed with the SEQUENASE<sup>R</sup> 2.0 system (United States Biochemical, Cleveland, OH, Tabor & Richardson, 1987). For automated DNA sequencing, the Tag DyeDeoxy<sup>TM</sup> Terminator Cycle Sequencing Kit (Applied Biosystems Inc., Foster City, CA) was utilized. Western blotting was performed following a modification of the method by Anderson *et al.* (1982, Electrophoresis 3:135). Sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting reagents were from Pierce (Rockford, IL). Polyclonal anti-rmuMIF antiserum was prepared from rabbits immunized with purified rmuMIF. Thioglycollate broth (Difco, Detroit, MI) was prepared according to the manufacturer's recommendation, autoclaved, and stored protected from light at room temperature. *E. coli* O111:B4 LPS and polymyxin B were purchased from Sigma (St. Louis, MO). LPS was resuspended in pyrogen-free water, vortexed, sonicated, aliquoted (5 mg/ml), and stored at -20°C. Serial dilutions of LPS were prepared in pyrogen-free water and sonicated (Branson 3210, Danbury, CT) for 10 min prior to use.

#### 6.1.2. MOLECULAR CLONING OF MURINE AND HUMAN MIF

Murine MIF was cloned from the mouse anterior pituitary cell line AtT-20/D16v-F2 (American Type Culture Collection, Rockville, MD). Cells were plated at  $1 \times 10^6$  cells/ml in DMEM containing 50 µg/ml gentamicin (Gibco BRL), and 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT). After 3 h of



incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, non-adherent cells were removed and the remaining adherent cells washed twice with DMEM/10% FBS. LPS (50 µg/ml) then was added and the cells were incubated  
5 for 16 h. At the end of this time, cell culture medium was removed and total RNA was isolated with RNAzol<sup>TM</sup>B according to the manufacturer's instructions. One µg of RNA was reverse transcribed using oligo (dT)<sub>12-18</sub> and M-MLV reverse transcriptase in a 50 µL  
10 reaction. Five µL of cDNA was amplified by PCR (32 cycles; 1 min at 94°C, 1 min at 55°C, 1 min at 72°C) using MIF primers. A single DNA amplification product of expected size was obtained and purified using the GENE CLEAN II<sup>R</sup> Kit (BIO 101 Inc., La Jolla, CA). The  
15 cDNA then was cloned into the plasmid pT7Blue and transformed into Nova Blue competent *E. coli* using the pT7Blue T-Vector Kit (Novagen, Madison, WI). Recombinant DNA was prepared from multiple white colonies using the Plasmid Magic<sup>TM</sup> Minipreps DNA  
20 Purification System (Promega) and sequenced manually in a Sequi-Gen II Sequencing Cell (BIO-RAD, Hercules, CA).

For human MIF cloning, Jurkat H33HJ-JA1 cells (American Type Culture Collection) were plated at 1 x  
25 10<sup>6</sup> cells/ml in RPMI containing 50 µg/ml gentamicin and 10% heat-inactivated FBS. After 3 h of incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, non-adherent cells were removed and the remaining adherent cells washed twice with RPMI/10% FBS. Cells were  
30 incubated for another hour and the total RNA was isolated and subjected to RT as described above. NdeI/MIF- and BamHI/MIF-fusion primers (5'-CCATATGCCGATGTTTCATCGTAAACAC-3' and 3'-CGGATCCTGCGGCTCTTAGGCGAAGG-5') were designed from a  
35 huMIF cDNA sequence (Weiser et al., 1989, Proc. Natl.

Acad. Sci. USA 86: 7522) and used to amplify human MIF cDNA as described above. A single PCR product of predicted size was obtained, purified using the GENE CLEAN II<sup>R</sup> Kit, and ligated into the NdeI/BamHI-digested pET11b vector (Novagen). *E. coli* DH5 $\alpha$  was transformed with the ligation mixture and the single recombinant colonies isolated after overnight growth. Plasmid DNA was prepared from eight clones and the MIF insert sequenced bi-directionally by automated methods using an ABI Model 373A DNA sequencer (Applied Biosystems Inc.).

#### 6.1.3. EXPRESSION AND PURIFICATION OF RECOMBINANT MIF

The recombinant pT7Blue clone containing muMIF cDNA was digested with the restriction enzymes NdeI and BamHI, and the MIF insert isolated and ligated into the NdeI/BamHI-digested pET11b vector (Novagen). *E. coli* DH5 $\alpha$  was transformed and the single recombinant colonies isolated and stored in 20% glycerol at -80°C until use. Murine or human MIF-containing pET11b plasmid DNA then was prepared and used to transform the *E. coli* BL21(DE3) expression strain (Novagen). One-liter cultures were grown at 37°C until the optical density at 600 nm reached 0.6-0.8. Isopropyl 1-thio- $\beta$ -D-galactopyranoside (IPTG) was added to a final concentration of 1 mM and the incubation continued at 37°C for an additional 3 h. Bacteria then were harvested by centrifugation and the cell pellets frozen at -20°C until use.

For protein purification, the bacterial pellets (corresponding to 50 ml of culture) were thawed and resuspended in 3.5 ml of Tris-buffered saline (50 mM Tris-HCl, 150 mM NaCl, pH 7.5). The bacteria were lysed by adding an equal volume of washed glass beads (106 microns; Sigma, G-8893) and vortexing the mixture

vigorously for 10 min. Lysis was confirmed by examination under phase contrast microscopy. Glass beads were removed by centrifugation at 1000 g for 10 min and the bacterial extract then was centrifuged at 38000 g for 30 min. The supernatant, representing the cleared bacterial lysate, was sterile-filtered through a 0.45  $\mu$ m and then a 0.22  $\mu$ m membrane filter and subjected to MONO Q anion exchange chromatography using a Fast Protein Liquid Chromatography system (FPLC) (Pharmacia, Piscataway, NJ). The Mono Q column was equilibrated with Tris-buffered saline. MIF was eluted with the same buffer in the first flow-through fractions (3ml), which were pooled and placed on ice immediately. The MIF-containing fractions then were applied to a C8-SepPak reverse-phase (RP) column (Waters, Division of MILLIPORE, Milford, MA) that had been washed first with methanol, followed by water. Unbound material was eluted by washing the resin with 10 column volumes of water and 20% acetonitrile/water, respectively. MIF then was eluted with 6 column volumes of 60% acetonitrile/water, frozen at -80°C, lyophilized, and kept at -20°C until use. For renaturation, MIF was dissolved at a concentration of 200-400  $\mu$ g/ml in 20 mM sodium phosphate buffer (pH 7.2) containing 8 M urea and 5 mM DTT, and dialyzed against 20 mM sodium phosphate buffer (pH 7.2) containing 5 mM DTT, followed by 20 mM sodium phosphate buffer (pH 7.2) alone. Renatured MIF was sterile-filtered and kept at 4°C until use. The LPS content of MIF preparations was determined by the chromogenic *Limulus* amoebocyte assay (Bio-Whittaker Inc., Walkersville, MD). Attempts to purify rmUMIF from the cleared bacterial lysate by affinity chromatography with S-hexyl-glutathione-agarose beads (Sigma, H-7011) were performed following the method

for single-step purification of protein/glutathione-S-transferase fusion constructs (Smith & Johnson, 1988, Gene 67: 31).

5                   6.1.4. PURIFICATION OF NATIVE MIF

Two grams of mouse liver acetone powder (Sigma, L-8254) were resuspended in 20 ml of Tris-buffered saline, vortexed, and the insoluble material removed by centrifugation at 1000 g for 10 min. The  
10 supernatant, containing MIF and other hepatic proteins, was filtered (0.45  $\mu$ m followed by 0.22  $\mu$ m filter) and subjected to MONO Q/FPLC anion exchange chromatography as described above. MIF eluted with the first flow-through fractions (3 ml), which were  
15 pooled and applied to a Pro S cation exchange column (BIO-RAD) that was equilibrated with Tris-buffered saline. MIF eluted with Tris-buffered saline and the MIF-containing fractions again were recovered from the flow-through. The MIF fractions then were pooled and  
20 applied to a C8-SepPak cartridge. MIF was eluted with 60% acetonitrile/water, lyophilized, and stored as described above. Attempts to purify liver MIF from mouse liver supernatant by affinity chromatography with S-hexyl-glutathione-agarose beads were performed  
25 as described above.

6.1.5. BIOCHEMICAL CHARACTERIZATION OF MIF

SDS-polyacrylamide gel electrophoresis (SDS-PAGE) was performed in 18% gels under reducing conditions  
30 (Laemmli, 1970, Nature 227: 680). The gels were either stained with silver or processed further for Western blotting. For silver staining, gels first were fixed for 16 h in 50% methanol/10% acetic acid and then analyzed as described (Poehling & Neuhooff,  
35 1981, Electrophoresis 2: 141). For Western blotting,

proteins were transferred to nitrocellulose membrane (Schleicher & Schuell, Keene, NH) by electroblotting at 50 V and 150 mA for 16 h using CAPS-transfer buffer (10 mM CAPS, 20% methanol, pH 11.0). Membranes then  
5 were incubated with blocking buffer (50 mM Tris-HCl, 500 mM NaCl, pH 7.5, 5% non-fat dry milk, 0.05% Tween-20) followed by incubation with a 1:1000 dilution of polyclonal rabbit anti-rmuMIF antiserum in binding buffer (50 mM Tris-HCl, 500 mM NaCl, pH 7.5, 1% BSA,  
10 0.05% Tween-20). After extensive washing in binding buffer, membranes were incubated with a 1:1000 dilution of horseradish peroxidase-conjugated goat anti-rabbit antibody (Pierce) in binding buffer, washed twice with binding buffer, twice with 50 mM  
15 Tris-Base (pH 7.5) containing 150 mM NaCl, and developed with chloronaphthol/H<sub>2</sub>O<sub>2</sub> substrate. Pre-stained protein molecular weight (M<sub>r</sub>) markers ranged from 2.5-43 kDa (Gibco BRL).

N-terminal MIF sequence was determined by Edman  
20 degradation (Allen, 1981 In: Sequencing of Proteins and Peptides, Elsevier, Amsterdam, New York) and automated gas-phase sequencing following a manufacturer's protocol (Applied Biosystems).

For N-glycosylation analysis, 0.1 µg of purified  
25 liver-derived MIF was incubated with the endoglycosidase PNGase F according to the manufacturer's instructions (New England BioLabs, Beverly, MA). Samples then were mixed 1:1 with Laemmli electrophoresis buffer, boiled for 5 min, and  
30 ¼ of the incubation mixture analyzed by SDS-PAGE/Western blotting as described above. To control for digestion efficiency, IgG light and heavy chains were digested in parallel and resolved on the same gel. Digestion was assessed to be complete after 3 h  
35 incubation with 5000 units/ml of PNGase F.

- For analytical size exclusion chromatography, 40  $\mu$ g of purified rmuMIF or rhuMIF was dissolved in 200  $\mu$ l 20 mM sodium phosphate buffer (pH 7.2) containing 7 M GdnHCl. The sample was separated over a Superose 12 HR 10/30 column (Pharmacia), equilibrated with the same buffer, and eluted at a flow rate of 0.5 ml/min. Size markers were from BIO-RAD (1.35-670 kDa) and were chromatographed under the same conditions.
- Mass spectrometric analysis (MS) of MIF was performed by matrix-assisted laser desorption ionization MS as described elsewhere (Hillenkamp & Karas, 1990, Meth. Enzymol. 193: 280) using a Shimadzu Kratos Kompact MALDI 3 V3.0.2 machine (Hannover, Germany). Twenty individual spectra were accumulated for each mass analysis. In addition, rmuMIF was analyzed by ion-spray mass spectroscopy (IS-MS) using an API III triple quadrupole mass spectrometer with an IonSpray<sup>TM</sup> interface (Sciex, Toronto).
- Purified, renatured rmuMIF (20  $\mu$ g/ml) and MIF-containing bacterial lysates (estimated MIF content: 50  $\mu$ g/ml) were analyzed photometrically for glutathione-S-transferase (GST) activity by the method of Fjellstedt et al. (1973, J. Biol. Chem. 248: 3702) using 1,2-epoxy-3-(p-nitrophenoxy)propane (EPNP) as a substrate and bovine GST (Sigma, G-8386) as a positive control (Blocki et al., 1992, Nature 360: 269). In this assay system, bovine GST (400  $\mu$ g/ml) was found to have an enzyme activity of 0.014 units/ml with respect to the substrate EPNP.

#### 6.1.6. BIOACTIVITY PROFILE

- The macrophage/monocyte migration inhibitory activity of recombinant MIF was analyzed by studying the migration of human peripheral blood monocytes in

modified Boyden chambers (Boyden, 1962, J. Exp. Med. 115: 453). Monocytes were isolated from heparinized venous blood of healthy donors and resuspended at a concentration of  $2.5 \times 10^6$  cells/ml in Gey's balanced salts solution (Gibco BRL) containing 2% fatty acid-free bovine serum albumin (Sigma) and 20 mM Hepes (pH 7.2). MIF-containing ( $0.0001$ - $10 \mu\text{g/ml}$ ) solutions or buffer controls were placed in the lower compartment of the Boyden chamber, covered tightly by a polyvinylpyrrolidone-free polycarbonate filter ( $5\text{-}\mu\text{m}$  pore size, no. 155845, Costar Corp., Cambridge, MA) and monocytes ( $1 \times 10^5$ ) added to the compartment above the filter. The chambers then were incubated for 3 h at  $37^\circ\text{C}$  in a humidified atmosphere with 5%  $\text{CO}_2$ . At the end of this time, the filters were recovered and the cells fixed and stained with Giemsa reagent. Monocytes then were counted as described previously (Sherry et al., 1992, Proc. Natl. Acad. Sci. USA 89: 3511).

TNF $\alpha$  production was quantitated in MIF-conditioned RAW 264.7 macrophage supernatants by L929 cell cytotoxicity as described previously (Wolpe et al., 1988, J. Exp. Med. 167: 570). RAW 264.7 macrophages were resuspended in RPMI/10% FBS, plated at  $1 \times 10^6$  cells/ml, incubated for 3 hours at  $37^\circ\text{C}$  in a humidified atmosphere with 5%  $\text{CO}_2$ , and washed twice with RPMI/1% FBS. Cells were incubated for 12-14 hours with various doses of MIF diluted in RPMI/1% FBS. At the end of each experiment, cell culture media were collected, centrifuged (10 min at 800 g), and supplemented with 1 mM PMSF. MIF-induced TNF $\alpha$  activity was measured immediately afterwards. Polyclonal rabbit anti-recombinant murine TNF $\alpha$  antiserum ( $50 \mu\text{l/ml}$ ) blocked TNF $\alpha$  activity completely, and MIF did not contribute to TNF $\alpha$  activity as

recombinant murine TNF $\alpha$  (rTNF $\alpha$ ) (5 pg/ml to 1  $\mu$ g/ml) cytotoxicity remained unchanged when rmuMIF (10 pg/ml to 10  $\mu$ g/ml) or anti-rmuMIF antibody were added to L929 cells together with rTNF $\alpha$ .

- 5 Nitric oxide (NO) production was quantitated in MIF-conditioned RAW 264.7 and thioglycollate-elicited peritoneal exudate macrophage supernatants by measuring nitrite and nitrate content with the Griess reagent. RAW 264.7 macrophages were prepared and
- 10 conditioned as described above. In some experiments, cells were incubated for 1 h with recombinant murine interferon- $\gamma$  (IFN- $\gamma$ ) (100 IU/ml) (Boehringer-Mannheim, Indianapolis, IN) prior to the addition of MIF. Thioglycollate-elicited peritoneal exudate macrophages
- 15 were obtained from BALB/c mice that were injected intra-peritoneally 3-4 days previously with 2 ml of sterile thioglycollate broth. Cells were harvested under strict aseptic conditions by lavage of the peritoneal cavity with 5 ml of an ice-chilled 11.6%
- 20 sucrose solution. After centrifugation (10 min at 800 g), cells were resuspended in RPMI/10% FBS and plated at a density of  $2 \times 10^6$  cells/ml. After 3 h of incubation, non-adherent cells (i.e. PMNs, lymphocytes) were removed with RPMI/1% FBS and the
- 25 remaining adherent cells conditioned with MIF as described for RAW 264.7 cells. Trace concentrations of contaminating LPS were neutralized by incubating MIF (1 and 10  $\mu$ g/ml in cell culture media) with polymyxin B at a concentration of 10 and 100 ng/ml,
- 30 respectively, for 30 min at room temperature under sterile conditions. The mixture was cleared by centrifugation and added to the prepared macrophage cultures. The polymyxin B concentration added was approximately a 1000-fold higher than necessary to
- 35 neutralize the contaminating LPS present. For



antibody neutralization of MIF activity, rmuMIF (1 and 10  $\mu\text{g/ml}$  in cell culture media) was treated with 20  $\mu\text{L/ml}$  of anti-rmuMIF or normal rabbit control serum and added to the macrophages as described above.

5

#### 6.1.7. CONFORMATIONAL AND STRUCTURAL STABILITY ANALYSIS

CD spectra were recorded on an Aviv Associates Model 62DS spectropolarimeter. The spectra represent the average of three scans recorded at 25°C in the range between 190 nm and 250 nm and were collected at 0.25 nm intervals, with a band width of 1.5 nm and a time constant of 1.0 sec. The quartz cells (1 and 10 mm) used in all CD measurements were washed with 30% HCl in ethanol, rinsed with water and methanol, and dried before used. Protein concentrations were determined from stock solutions prepared in 20 mM phosphate buffer (pH 7.2) using the Bio-Rad protein assay (BIO-RAD). This assay was found to agree with values obtained by quantitative amino acid analysis. Thirty min before CD analysis, proteins were diluted from the stock solutions to a final concentration of 10  $\mu\text{M}$  in 20 mM phosphate buffer (pH 7.2), unless stated otherwise. CD spectra are presented as a plot of the mean molar ellipticity per residue ( $[\theta]$ ,  $\text{deg cm}^2 \text{dmol}^{-1}$ ) versus the wavelength.

For unfolding experiments, MIF stock solutions dissolved in 20 mM phosphate buffer (pH 7.2) were mixed with increasing volumes of 8 M GdnHCl (molecular biology reagent grade, Sigma) prepared in 20 mM phosphate buffer (pH 7.2) so as to achieve a final protein concentration of 10  $\mu\text{M}$  and the indicated final GdnHCl concentrations. Samples then were equilibrated at room temperature for 30 min prior to the recording of CD spectra.

## 6.2. RESULTS

### 6.2.1. CLONING OF MURINE AND HUMAN MIF

N-terminal protein sequencing and cDNA cloning  
5 were used to identify the murine homolog of MIF among  
the proteins secreted by LPS-stimulated anterior  
pituitary cells. MuMIF cDNA was prepared from the  
total RNA of LPS-stimulated anterior pituitary cells  
(AtT-20) and amplified with MIF primers. MIF cDNA  
10 then was cloned into the pT7Blue T-vector and  
subjected to DNA sequence analysis. The muMIF cDNA  
sequence obtained from 6 plasmid clones was compared  
to a previously published human T-cell MIF cDNA  
(Weiser et al., 1989, Proc. Natl. Acad. Sci. USA 86:  
15 7522). Murine MIF cDNA (FIG. 3) was found to display  
a 88.2% sequence homology to this huMIF sequence over  
a 348 nucleotide open reading frame and to be  
identical in sequence with recently reported murine  
3T3 fibroblast, and murine embryonic eye lens MIF  
20 cDNAs (Lanahan et al., 1992, Mol. Cell. Biol. 12:  
3919; Wistow et al., 1993, Proc. Natl. Acad. Sci. USA  
90: 1272).

HuMIF cDNA was prepared by RT/PCR of RNA isolated  
from resting Jurkat H33HJ-JA1 T-cells. MIF cDNA was  
25 amplified with flanking primers bearing NdeI/BamHI  
restriction sites, thus enabling the subsequent  
cloning of the huMIF amplification product directly  
into a NdeI/BamHI-digested pET11b prokaryotic  
expression plasmid. Human MIF cDNA sequence then was  
30 obtained by sequencing both DNA strands of 8  
independently derived *E. coli* clones. This Jurkat T-  
cell MIF sequence was found to be identical to a  
recently reported human fetal lens MIF cDNA (Wistow et  
al., 1993, Proc. Natl. Acad. Sci. USA 90: 1272) and to  
35 a glycosylation inhibition factor (GIF) cDNA (Mikayama

et al., 1993, Proc. Natl. Acad. Sci. USA 90: 10056), but differed from the first reported human T-cell MIF sequence obtained from the T-CEMB cell line (Weiser et al., 1989, Proc. Natl. Acad. Sci. USA 86: 7522) by  
5 a single base change at position 318. This G → A substitution produces a conservative Ser<sup>106</sup> → Asn<sup>106</sup> change in the deduced amino acid sequence and renders the human protein even more homologous with murine MIF. Murine pituitary MIF cDNA and human Jurkat cell  
10 cDNA exhibited 88.5% identity over a 348 nucleotide open reading frame and the amino acid sequences deduced for murine (AtT-20) and human (Jurkat) MIF were found to be 90% identical over the 115 amino acids (FIG. 4). No apparent N-terminal signal  
15 sequences were evident in either the murine or the human proteins. MIF thus joins a growing list of cytokines, such as interleukin-1 (IL-1) (Rubartelli et al., 1990, EMBO J. 9:1503), basic fibroblast growth factor (bFGF) (Jackson et al., 1992, Proc. Natl. Acad.  
20 Sci. USA 89: 10691), and a secreted form of cyclophilin (Sherry et al., 1992, Proc. Natl. Acad. Sci. USA 89: 3511), which are released from cells by non-classical protein secretion pathways. Two potential N-glycosylation sites were found at  
25 positions 75 and 110 of the muMIF amino acid sequence and were at nearly identical positions in the human Jurkat MIF sequence (positions 73 and 110). Three cysteine residues (positions 57, 60, and 81) also were at identical places in the murine and human MIF  
30 predicted amino acid sequences.

#### 6.2.2. EXPRESSION AND PURIFICATION OF RECOMBINANT MIF FROM *E. COLI*

Recombinant muMIF and huMIF were expressed in  
35 *E. coli* by cloning murine and human MIF cDNA into the IPTG-inducible pET expression plasmid system. Initial

attempts to express murine MIF from the pET15b vector, which created an N-terminal oligo-histidine fusion protein and allowed for facile purification of recombinant protein by ion metal affinity chromatography (IMAC), were unsuccessful because the protein was resistant to the subsequent thrombin cleavage necessary to remove the oligo-histidine leader. Thus, recombinant muMIF was expressed and then purified by conventional means and the muMIF cDNA subcloned into the plasmid pET11b. This produced a construct which bore a three amino acid (Met-Asp-Ser) leader sequence attached to the N-terminus of MIF. HuMIF also was expressed from the pET11b vector but was engineered by DNA amplification to begin at the second amino acid of the open reading frame, with the start methionine of MIF provided by the NdeI restriction site. The correct expression of MIF was verified by N-terminal amino acid sequencing of gel-purified protein (10 amino acids for muMIF), SDS-PAGE, and Western blotting with anti-AtT-20 MIF antibody. The pET11b-derived MIF was used for the subsequent purification of bioactive muMIF and huMIF.

Recombinant murine or human MIF were found to constitute 40% of the total supernatant protein of *E. coli* lysates. Anion exchange chromatography at pH 7.5 resulted in approximately an 80% purification of murine and human MIF. Subsequent application to a C8 reverse-phase column yielded pure protein for both the murine and the human recombinant proteins, as verified by the appearance of single bands of predicted  $M_r$  (12.5 kDa) by SDS-PAGE/silver staining and by mass spectrometric analysis, *infra*. This simple and rapid two-step procedure was used to purify 1 mg of murine or human MIF per 50 ml of *E. coli* culture. After renaturation, 0.8 - 0.9 mg of soluble, bioactive

recombinant MIF was obtained per 50 ml culture. The overall yield of MIF protein from total bacterial supernatant protein was estimated to be 20%. This two-step purification method was elected for reasons of simplicity and rapidity and to minimize protein losses by non-specific precipitation. Purification by C8 chromatography using a disposable, low volume push column also served to remove the LPS carried over from *E. coli* host cells. Recombinant MIF purified by these procedures contained no more than trace amounts of LPS (4-8 ng LPS per mg MIF).

#### 6.2.3. PURIFICATION OF NATIVE MIF

To assess more precisely the biological activities of recombinant MIF, native MIF was also purified from mouse tissue. Pituitary cells yielded insufficient quantities of MIF protein for biochemical characterization, but Western blotting showed that the liver was an abundant source of MIF *in vivo*. MIF was purified from liver by a method similar to that used for recombinant MIF purification. Mono Q anion exchange chromatography resulted in partial purification and increased the relative MIF content by approximately 25-fold. Subsequent application to the C8 push column afforded almost pure MIF. Upon SDS-PAGE/silver staining analysis however, 5 additional bands were detected that could not be removed by varying the elution conditions of the RP chromatography. A Pro S cation exchange chromatography step therefore was added prior to the C8 step to remove these contaminants. The purity and homogeneity of liver MIF was established by SDS-PAGE/silver staining, Western blotting using anti-rmuMIF antibody, and mass spectrometric analysis. Laser desorption MS demonstrated a single protein

species of  $M_r$  12,555. The MIF content of mouse liver acetone powder was estimated to be less than 0.1% of total protein and 2 grams of mouse liver powder were used to purify to homogeneity 50  $\mu$ g native liver MIF.

5

#### 6.2.4. BIOCHEMICAL CHARACTERIZATION OF MIF

The  $M_r$  for liver murine MIF, rmuMIF, and rhuMIF were each between 12000 and 13000 as estimated by reducing SDS-PAGE. Two potential N-linked  
10 glycosylation sites were detected in the predicted amino acid sequence of murine and human MIF and thus the possibility that native muMIF was post-translationally glycosylated *in vivo* was investigated. Endoglycosidase F (PNGase F) digestion of purified  
15 liver MIF followed by reducing SDS-PAGE/Western blotting analysis showed no shift in the observed  $M_r$ , indicating the absence of significant N-glycosylation of native liver MIF.

Gel exclusion chromatography of GdnHCl-denatured  
20 recombinant MIF showed a  $M_r$  of 14,256 and 13,803 for the murine and human proteins, respectively, indicating that MIF eluted predominantly as a monomer.

Laser desorption MS of MIF was performed to further assess glycosylation as well as the presence  
25 of any other significant post-translational modifications of MIF. Native MIF obtained from mouse liver was determined to have a  $M_r$  of 12,555 (predicted  $MH^+$  average mass of oxidized muMIF = 12,503). By comparison, rmuMIF (which bore a 3 amino acid N-terminal leader sequence) was found to have a  $M_r$  of  
30 12,814 (predicted  $MH^+$  average mass of oxidized rmuMIF = 12,801). Ion-spray MS of rmuMIF yielded a similar mass ( $M_r$  = 12,804). Using laser desorption MS, rhuMIF was found to have a  $M_r$  of 12,521 (predicted  $MH^+$  average  
35 mass of oxidized rhuMIF = 12,475). Differences

observed between the experimental and theoretical, predicted masses were within the margin of experimental error (0.1-1%) for these analyses. These data essentially ruled out the presence of significant post-translational modifications of either native or recombinant MIF protein.

A 12 kDa protein purified from rat liver bearing N-terminal homology with huMIF has been reported to bind to glutathione affinity matrix and to exhibit glutathione-S-transferase (GST) activity (Blocki et al., 1992, Nature 360: 269). Neither liver MIF nor rmuMIF was shown to specifically absorb to glutathione-modified matrix.

#### 6.2.5. BIOACTIVITY PROFILE

Purified rmuMIF displayed significant migration inhibitory activity when tested on human peripheral blood monocytes (FIG. 5). A bell-shaped dose response curve was observed with peak activity occurring at 0.1 µg/ml. The precise basis for diminished inhibitory activity at high MIF concentrations is not known, but similar dose-response profiles have been observed for chemotactic cytokines (Sherry et al., 1992, Proc. Natl. Acad. Sci. USA 89: 3511).

Recombinant MIF was used to induce TNFα release by the murine RAW 264.7 macrophage cell line. Both rmuMIF and rhuMIF added at concentrations of 0.1-10 µg/ml caused the release of bioactive TNFα in the ng/ml range (Table II). Despite species-specific differences, rhuMIF at 1/µg/ml was found to be slightly more active than rmuMIF on murine macrophages. At the same concentration (1 µg/ml), the TNFα-inducing activity of MIF obtained from mouse liver was found to be higher than the activity of rmuMIF. When tested at a concentration of 0.1 µg/ml

however, the TNF $\alpha$ -inducing activity of mouse liver MIF was identical to rmuMIF. Similar findings were obtained when thioglycollate-elicited mouse peritoneal macrophages were used to study MIF-induced TNF $\alpha$

5 release. Overall, the activity of purified native MIF was within the range observed for the recombinant proteins.

**TABLE II**

10 **MIF-INDUCED SECRETION OF TNF $\alpha$  BY MOUSE MACROPHAGES\***

Secretion of TNF $\alpha$ (ng/ml) induced by:			
MIF ( $\mu$ g/ml)	rmuMIF	rhuMIF	Native MIF
15 10	ND	3.9 $\pm$ 2.8	ND
1	1.2 $\pm$ 0.1	3.4 $\pm$ 2.8	6.9 $\pm$ 3.4
0.1	0.2 $\pm$ 0.3	2.9 $\pm$ 2.4	0.2 $\pm$ 0.1
0.01	0	ND	0

20 \*RAW 264.7 macrophages were prepared and incubated with MIF for 12-14 h as described in Section 6.1.6. supra. After the incubations, macrophage culture supernatants were removed, and analyzed by L929 cell toxicity assay for TNF $\alpha$  activity. TNF $\alpha$  measurements were performed in duplicate and TNF $\alpha$  activity is  
 25 expressed as the difference between the level produced by MIF-stimulated cells and by non-stimulated control cells. Data are the mean $\pm$ SEM of at least three separate experiments. ND: not determined.

Native and recombinant MIF also induced NO production in macrophages (Table III). However,  
 30 significant MIF-induced NO production was observed only when macrophages were incubated with IFN- $\gamma$  prior to stimulation with MIF. In RAW 264.7 macrophages, the activity of rhuMIF at 1  $\mu$ g/ml was found to be approximately 4-fold higher than rmuMIF. Native MIF  
 35 obtained from mouse liver also was more active in this



assay than rmuMIF. MuMIF stimulated NO production from IFN- $\gamma$ -primed, thioglycollate-elicited peritoneal exudate macrophages at levels which were comparable with those observed in RAW 264.7 cells. However,  
5 liver MIF at 1  $\mu$ g/ml was not found to be more active than rmuMIF on this macrophage cell type.

The synthesis of NO by macrophages is induced by  $\geq 10$  ng/ml of LPS and this effect is potentiated by IFN- $\gamma$  priming (Ding et al., 1988, J. Immunol. 141:  
10 2407). To exclude the possibility that the trace amounts of LPS present in recombinant MIF preparations contribute to NO production, MIF-induced NO release was tested after neutralization of LPS with a large excess of polymyxin B. The production of NO was not  
15 reduced by polymyxin B treatment, arguing against a potentiating role for LPS in the stimulation of macrophages by MIF. In further control studies, the addition of neutralizing anti-rmuMIF antiserum (20  $\mu$ l/ml) inhibited rmuMIF (1 and 10  $\mu$ g/ml)-induced  
20 NO production, further confirming the specificity of macrophage stimulation by recombinant MIF.

25

30

35

**TABLE III**  
**NO PRODUCTION IN MOUSE MACROPHAGES INDUCED**  
**BY MIF, ALONE OR IN COMBINATION WITH IFN- $\gamma$ <sup>a</sup>**

5	Formation of nitrite ( $\mu$ M) induced by:				
		MIF ( $\mu$ g/ml)	rmu MIF	rhuMIF	Native MIF
	- IFN- $\gamma$	10 <sup>a</sup>	0.2 $\pm$ 0.2	3.7 $\pm$ 2.1	ND
		1 <sup>a</sup>	0.5 $\pm$ 0.5	1.5 $\pm$ 1.0	0
10		0.1 <sup>a</sup>	0.3 $\pm$ 0.3	0	0
		0.01 <sup>a</sup>	0	ND	0
	+ IFN- $\gamma$	10 <sup>a</sup>	12.3 $\pm$ 2.5	25.3 $\pm$ 7.5	ND
		1 <sup>a</sup>	5.7 $\pm$ 1.3	20.4 $\pm$ 5.7	9.8 $\pm$ 1.3
15		0.1 <sup>a</sup>	1.7 $\pm$ 0.7	8.4 $\pm$ 2.6	3.1 $\pm$ 1.0
		0.01 <sup>a</sup>	1.9 $\pm$ 1.2	ND	0.5 $\pm$ 0.2
		1 <sup>b</sup>	12.5	ND	3.5

<sup>a</sup>RAW 264.7 macrophages were prepared and incubated with MIF for 12-14 h. At the end of incubation, 300  $\mu$ L of culture supernatant was removed, mixed with 600  $\mu$ L of Griess reagent, and the concentration of nitrite measured. Nitrite measurements were performed in duplicate and nitrite production is expressed as the difference between the level produced by MIF-stimulated cells versus non-stimulated, or IFN- $\gamma$ -treated control cells. Data are the mean $\pm$ SEM of at least three separate experiments. ND: not determined.

<sup>b</sup>NO production by MIF-stimulated thioglycollate-elicited peritoneal macrophages in place of RAW 264.7 macrophages. Data are the mean of two separate experiments.

#### 6.2.6. CONFORMATIONAL AND STRUCTURAL STABILITY ANALYSES

Far-UV CD has been used widely to analyze the solution conformation of proteins and to verify native and renatured protein structures (Greenfield & Fasman, 1969, Biochemistry 8: 4108). Far-UV CD spectroscopy

and GdnHCl-induced unfolding studies were performed with recombinant MIF to begin to assess the secondary structure and the structural stability of this protein in a solution environment. No concentration dependent

5 changes in the CD spectra of MIF were observed within the range studied (1.6-32  $\mu$ M MIF). The spectrum for rmuMIF showed a pronounced positive ellipticity at 192 nm, a broad negative ellipticity between 209 nm and 222 nm, and small distinct minima at 210 nm and 222

10 nm. These data are consistent with a highly ordered secondary structure containing predominantly  $\beta$ -sheet, but also  $\alpha$ -helix conformation. The spectrum for rhuMIF showed a prominent positive ellipticity at 197 nm and a strong negative ellipticity between 211 nm

15 and 225 nm was observed. This CD spectrum also was consistent with a highly ordered secondary structure, but was suggestive of a higher  $\beta$ -sheet content and a lower  $\alpha$ -helix content than rmuMIF. Secondary structure compositions for both proteins then were

20 estimated from computational fits of the CD spectra over the 190-250 nm range using the method of Brahms & Brahms (1980, J. Mol. Biol. 138: 149), which is frequently employed for the analysis of  $\beta$ -sheet-rich proteins. Recombinant muMIF was estimated to contain

25 42.2%  $\beta$ -sheet, 15.3%  $\alpha$ -helix, 7.2%  $\beta$ -turn, and a remainder of 35.3% in random coil. Recombinant huMIF contained a somewhat greater extent of ordered structure, with 72.9%  $\beta$ -sheet, 0%  $\alpha$ -helix, 8.3%  $\beta$ -turn, and 17.3% random coil conformation. Secondary

30 structure calculations following the methods of Chou & Fasman (1978, Adv. Enzymol. 47: 45) and Garnier et al. (1978, J. Mol. Biol. 120: 97) also predicted that both muMIF and huMIF exhibit a highly ordered secondary structure. Chou/Fasman calculations predicted the

35 following secondary structure compositions- rmuMIF:

35.6%  $\beta$ -sheet, 10.6%  $\alpha$ -helix, and 21.2%  $\beta$ -turn;  
rhuMIF: 26.1%  $\beta$ -sheet, 30.4%  $\alpha$ -helix, and 17.4%  $\beta$ -  
turn. Garnier/Osguthorpe/Robson analysis predicted  
the following structures- rmuMIF: 39%  $\beta$ -sheet, 8.5%  $\alpha$ -  
5 helix, and 20.3%  $\beta$ -turn; rhuMIF: 24.4%  $\beta$ -sheet, 19.1%  
 $\alpha$ -helix, and 25.2%  $\beta$ -turn. The CD data, together with  
the secondary structure predictions according to the  
Chou/Fasman and Garnier/Osguthorpe/Robson  
calculations, indicate that both recombinant murine  
10 and human MIF contain a high fraction of  $\beta$ -sheet  
conformation (24.4% - 72.9%). The spectroscopic data  
and primary sequence predictions showed a similar  
 $\alpha$ -helical content for rmuMIF (8.5% - 15.3%). Although  
the secondary structures predicted by Chou/Fasman and  
15 Garnier/Osguthorpe/Robson calculations suggested that  
there was a significant  $\alpha$ -helical content in huMIF,  
the CD analysis did not show any detectable  $\alpha$ -helical  
elements in rhuMIF.

Helical conformations induced at the membrane-  
20 water interface have been suggested to be important  
for ligand membrane interactions and to influence the  
binding of ligands to receptors (Kaiser & Kezdy, 1983,  
Proc. Natl. Acad. Sci. USA 80:1137; Fry et al., 1992,  
Biopolymers 32:649; Erne et al., 1985, Biochemistry  
25 24: 4263). 2,2,2-Trifluoro-ethanol (TFE) induces and  
stabilizes the helical conformation of proteins with a  
helix-forming propensity and has been used to mimic  
the influence of membranes on polypeptide conformation  
(Sönnichsen et al., 1992, Biochemistry 31: 8790). To  
30 begin to test whether the  $\alpha$ -helical content of either  
murine or human MIF might be increased in the  
membranous environment, far-UV CD analysis of rmuMIF  
and rhuMIF was performed in 50% TFE. TFE  
significantly increased the  $\alpha$ -helical content of  
35 rmuMIF, as demonstrated by a larger positive

ellipticity at 192 nm and pronounced negative ellipticities at 208 nm and 222 nm. The calculated fraction of  $\alpha$ -helix conformation increased from 15.3% to 34.7% in the presence of TFE. Recombinant huMIF, which was found to have no measurable  $\alpha$ -helical elements in aqueous solution, showed a significant content of  $\alpha$ -helix in the presence of TFE as evident by a shift of the positive ellipticity from 197 nm to 192 nm and distinct negative ellipticities at 208 nm and 222 nm. Computational analysis of the spectroscopic data showed that the percentage of  $\alpha$ -helix increased from 0% to 36.3%. These data indicate that both murine and human MIF are likely to adopt significant  $\alpha$ -helical conformation in a membranous environment.

Protein structural stability can be quantitated by determining the free energy of unfolding,  $\Delta G_{N-U}$ , where N is the fraction of protein in the native state and U is the fraction in the unfolded state. One method used frequently to assess protein stability is to measure protein mean molar ellipticity per residue as a function of wavelength and GdnHCl concentration (Pace, 1975, CRC Crit. Rev. Biochem. 3: 1). Unfolding curves expressed as the percentage of unfolded protein relative to native protein (i.e. the change in ellipticity at 222 nm) over the concentration of GdnHCl provide two measures of structural stability: 1) the midpoint of unfolding,  $[GdnHCl]_{0.5}$ , which can be deduced from the linear part of the unfolding curve and 2) the free energy of unfolding at zero denaturant concentration,  $\Delta G_{N-U}^{\circ}$ , which can be extrapolated from the unfolding curve by the linear extrapolation method (LEM) (Santoro & Bolen, 1992, Biochem. 31: 4901). Both measures are based on the premise that unfolding follows a reversible two-state mechanism and that

unfolding free energy is linearly dependent on denaturant concentration (Greene & Pace, 1974, J. Biol. Chem. 249: 5388).

The CD spectra of rmuMIF were recorded between  
5 the wavelengths 210 nm - 250 nm in the presence of increasing concentrations of GdnHCl (0-7 M). Recording below 210 nm was not performed due to strong scattering effects at these wavelengths. The MIF spectrum, defined in the absence of GdnHCl, changed  
10 visibly when  $\geq 1.5$  M GdnHCl was present, indicating a significant loss of conformational integrity. The featureless spectrum for fully denatured MIF in the presence of 7 M GdnHCl was similar to spectra observed for other fully unfolded proteins (Greenfield &  
15 Fasman, 1969, Biochem. 8: 4108). For quantitation of GdnHCl-induced unfolding, the unfolding of MIF  $\alpha$ -helical structures was plotted by expressing the relative change in  $[\theta]$  at 222 nm (percent unfolded) with respect to increasing concentrations of GdnHCl.  
20 The midpoint of unfolding was observed to be at 1.8 M GdnHCl. Next, the LEM was applied to extrapolate  $\Delta G_{N-U}^{\circ}$  from a plot of  $\Delta G_{N-U}$  versus the concentration of GdnHCl( $[GdnHCl]$ ). Data points of the linear portion of the curve could be replotted according to the  
25 following equation:

$$\Delta G_{N-U} = \Delta G_{N-U}^{\circ} - m [GdnHCl]$$

where  $m$  is the slope of the curve, and subjected to linear regression analysis. Murine MIF thus was calculated to have a  $\Delta G_{N-U}^{\circ}$  of 11.75 kJ/mol. By  
30 comparison, globular proteins generally exhibit  $\Delta G_{N-U}^{\circ}$  values of approximately  $50 \pm 15$  kJ/mol (Pace 1990, Trends Biochem. Sci. 15: 14).

7. EXAMPLE: IN VIVO ADMINISTRATION OF MIF  
INCREASES ENDOTOXIN-INDUCED  
LETHALITY AND ANTI-MIF  
TREATMENT CONFERS PROTECTION

The following experiment was initially carried  
5 out to test the efficacy of purified MIF as a  
protective factor against endotoxin-induced lethality  
in animals. Surprisingly, the in vivo administration  
of MIF potentiated LPS lethality in mice and  
exacerbated the response to endotoxin challenge.  
10 Furthermore, mice pretreated with an anti-MIF  
antiserum were significantly protected against  
endotoxin-induced lethality. Anti-MIF treatment also  
protected against TNF $\alpha$ -induced lethality.

15 7.1. MATERIALS AND METHODS

7.1.1. MICE

Nine-week-old BALB/c mice (19-21 g of weight)  
were utilized for this study. A total of 20 animals  
20 in each category (i.e., LPS, LPS plus MIF, anti-MIF,  
preimmune serum and saline-treated) were included in  
the experiments described below.

7.1.2. IN VIVO TREATMENT WITH ENDOTOXIN AND MIF

25 Mice were injected i.p. with 15 mg/kg *E. coli*  
0111:B4 LPS (lipopolysaccharide) in saline. MIF  
prepared as described in Section 6, *supra*, was  
administered i.p. to mice at 5 mg/kg in saline at  
0 and 12 hours after LPS treatment. Both control and  
30 treated animal groups were maintained under the  
standardized conditions of the animal facility colony  
rooms and monitored for survival.

### 7.1.3. ANTI-MIF TREATMENT

Anti-MIF antiserum was produced by immunization of rabbits with 100  $\mu$ g of recombinant murine MIF purified from *E. coli* transfected with the coding  
5 sequence in an expression vector. Anti-MIF serum and normal rabbit serum (NRS) showed no cross-reactivity with *E. coli* 0111:84 LPS as assessed by Western blot analysis and LPS-specific ELISA.

Mice were pretreated with anti-MIF antiserum  
10 (containing < 0.2 ng LPS/ml), preimmune NRS (containing < 3 ng LPS/ml), or a saline control 24 hours and 2 hours prior to challenge with LPS (i.p. at 17.5 mg/ml, an amount of *E. coli* 011.B LPS that had been established as an LD<sub>75</sub> for saline-injected control  
15 mice in prior dose-ranging studies). Pretreatments consisted of 200  $\mu$ l of sera or 200  $\mu$ l 0.9% (w/v) NaCl as control, administered i.p.

TNF $\alpha$  challenge consisted of 0.7  $\mu$ g in Tris-buffered saline administered i.p. concurrently with 18  
20 mg D-galactosamine.

### 7.2. RESULTS

The effects of MIF on endotoxin-induced lethality were tested by comparing the survival curves of  
25 animals challenged in the presence and absence of administered exogenous recombinant MIF. Briefly, each animal was injected i.p. with 300  $\mu$ g LPS. Half of the animals were additionally administered 100  $\mu$ g MIF i.p. at 0 and 12 hours post-LPS challenge.

30 The pooled data representing the results of two such experiments are depicted in FIG. 6A in graphic form. These experiments demonstrate that MIF potentiated LPS lethality in the mice tested. First, the LPS/MIF-treated mice showed an accentuated rate of  
35 mortality in that a differential effect of MIF



administration, relative to that of the LPS-only mice, was seen beginning approximately 20 hours post LPS challenge. Second, the overall survival of animals in the two categories demonstrated the increased

5 cumulative mortality of animals exposed to MIF in addition to LPS, with only 15% of the LPS/MIF mice surviving at 84 hours post-LPS challenge compared to 65% of the LPS-only treated mice surviving to this time point. The overall comparison in a two tailed  
10 Fisher's exact test is  $P=0.003$ .

The effects of MIF on endotoxin- and  $\text{TNF}\alpha$ -induced lethality were tested by comparing the survival of animals challenged after pretreatment with anti-MIF antisera to those animals pretreated with preimmune  
15 sera and saline controls. Briefly, animals were injected, i.p., with either anti-MIF antisera, preimmune sera, or saline, 24 hours and 2 hours prior to challenge with either LPS or  $\text{TNF}\alpha$ .

FIG. 6B illustrates the pooled data representing  
20 the results of two experiments in which the mice were challenged with LPS. As can be seen, mice pretreated with anti-MIF antisera were significantly protected from LPS-induced lethality, with all mice surviving until the last recorded timepoint, 86 hours post-  
25 challenge. In contrast, only 25% of the animals that had been pretreated with the saline control were still alive at 84 hours post-LPS challenge, and only 50% of those pretreated with preimmune sera were alive. A statistical analysis of these results indicated that  
30 the chances of these results occurring under the null hypothesis were, for the overall comparison:  
 $P<0.00001$ ; anti-MIF vs. saline :  $P<0.00001$ ; anti-MIF vs. NRS : 0.0004; and NRS vs. saline : 0.19.

Table IV illustrates the pooled data representing  
35 the results of three experiments in which the mice

were challenged with TNF $\alpha$  and galactosamine. As recorded above in similar experiments with LPS challenge, mice pretreated with anti-MIF antisera were significantly protected from TNF $\alpha$ -induced lethality, with all mice surviving until the last recorded timepoint. In contrast, only 50% of the animals that had been pretreated with the saline control were still alive at 120 hours post-TNF $\alpha$  challenge, and only 60% of those pretreated with preimmune sera were alive. Any marginal benefits afforded by pretreatment with preimmune control sera in these experiments most likely reflects the well-known but small beneficial effect of exogenous gamma globulin treatment.

---

**TABLE IV**

Treatment Group	Days after TNF $\alpha$ and galactosamine challenge				
	1	2	3	4	5
Saline	9/10 90%	9/10 90%	9/10 90%	5/10 50%	5/10 50%
Pre-immune	10/10 100%	10/10 100%	10/10 100%	6/10 60%	6/10 60%
Anti-MIF	10/10 100%	10/10 100%	10/10 100%	10/10 100%	10/10 100%

Fraction of the treatment groups surviving after TNF $\alpha$  + galactosamine treatment on day 0.

---

These experiments demonstrate that the immunological neutralization of MIF had the ability to protect against both LPS-induced and TNF $\alpha$ -induced lethality in the mice tested, as evidenced by the sparing by anti-MIF treatment of animals that would otherwise be expected to die. Thus, these results not

only prove that exogenously administered MIF has a deleterious effect on the shock process, but also demonstrate that the neutralization of endogenous MIF protects against endotoxin-induced and TNF $\alpha$ -induced lethality due to shock.

8. EXAMPLE: ADMINISTRATION OF EXOTOXIN  
INDUCES MIF SECRETION AND ANTI-MIF  
TREATMENT CONFERS PROTECTION

10

8.1. MATERIALS AND METHODS

8.1.1. CELL PREPARATION AND MIF INDUCTION

RAW 264.7 murine macrophages were obtained from the American Type Culture Collection (Rockville, MD). Cells were grown in RPMI 1640 medium (Gibco, Grand Island, NY), 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT), 50  $\mu$ g/ml of carbenicillin and gentamicin. RAW 264.7 macrophages were washed with fresh medium, harvested by gentle scraping, resuspended in RPMI/10%FBS, and incubated at  $3 \times 10^6$  cells/well in 3.5-cm tissue culture plates (Linbro\*, Flow, McLean, VA). After 3h of incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, nonadherent cells were removed and wells were washed twice with RPMI/1%FBS. Cells then were incubated for 12-15h with TSST-1 (at concentrations ranging from 100 ng/ml to 100 attog/ml). At the end of experiment, cell-culture media were collected and centrifuged (10 min at 800 g). Cell-conditioned media were concentrated 10-fold by membrane filtration (10 kD cut-off) (Centricon-10, Amicon, Beverly, NY).

Thioglycollate-elicited peritoneal macrophages were obtained from BALB/c mice that were injected i.p. 3-4 days previously with 2 ml of sterile thioglycollate broth. Cells were harvested under

strict aseptic conditions by lavage of the peritoneal cavity with 5 ml of an ice-chilled 11.6% sucrose solution. After centrifugation (10 min at 800 g), cells were resuspended in RPMI/10% FBS, enumerated and  
5 plated at a density of  $3-4 \times 10^6$  cells/well. Cells were incubated for 12 to 15h with TSST-1 (at concentrations ranging from 100 ng/ml to 100 attog/ml).

Murine T-cells (LBRM-33, a T lymphoma cell-line) were obtained from the American Type Culture  
10 Collection (Rockville, MD). Cells were grown in RPMI 1640 medium (Gibco, Grand Island, NY), 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT), 0.05 mM 2-mercaptoethanol and 50  $\mu$ g/ml of gentamicin. Cells were washed with fresh medium,  
15 harvested by gentle scraping, resuspended, and incubated at  $1 \times 10^6$  cells/well in 3.5-cm tissue culture plates (Linbro<sup>®</sup>, Flow, McLean, VA). After 3h of incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, nonadherent cells were removed and wells were  
20 washed twice with RPMI/1%FBS. Cells then were incubated for 12-15h with TSST-1 (at concentrations ranging from 100 ng/ml to 100 attog/ml). At the end of experiment, cell-culture media were collected and centrifuged (10 min at 800 g). Cell-conditioned media  
25 were concentrated 10-fold by membrane filtration (10 kD cut-off) (Centricon-10, Amicon, Beverly, NY).

Samples were resolved on 18% SDS polyacrilamide gels and analyzed by Western blotting as described above.

30

#### 8.1.2. ANTI-MIF TREATMENT

Eight to 10-week-old (19-21 g) female BALB/c were purchased from Charles River (Kingston, NY). Animals were housed in groups of 5 or 10 mice per cage with  
35 free access to food and water and were acclimatized

for 5 days prior to experimentation. BALB/C mice were injected i.p. with 200  $\mu$ l of polyclonal rabbit anti-MIF serum or normal rabbit serum (controls) followed 2 hr later by a lethal i.p. injection combining 50  $\mu$ g of TSST-1 with 20 mg D-galactosamine. Mice were observed for 7 days.

## 8.2. RESULTS

TSST-1 was found to be an extremely potent inducer of MIF secretion by macrophages (both RAW 264.7 and peritoneal macrophages) and T cells *in vitro*. Dose-response studies of TSST-1-induced MIF secretion showed two peaks of MIF secretion. The first peak occurred at 10 pg/ml of TSST-1 and the second at 100 attog/ml ( $=10^{-4}$ pg/ml). Anti-MIF treatment markedly reduced mortality from 87.5% (controls) to 33% ( $p=0.05$ , two-tailed Fisher's exact test) (FIG. 7).

## 9. EXAMPLE: MIF IS A PRIMARY INFLAMMATORY MEDIATOR IN PARASITIC INFECTIONS

### 9.1. PRODUCTION OF MIF IN MALARIA INFECTION

During malaria infection, hemozoin results from the pericyte-specific polymerization of heme molecules liberated by the catabolism of red blood cell hemoglobin by intra-erythrocytic forms of *Plasmodium*, and this "malaria pigment" accumulates within the reticuloendothelial system of infected individuals. The structure of hemozoin is known, and hemozoin can be chemically synthesized by polymerizing ferric heme in acetic acid, and separated from unreacted heme by sequential extraction in  $\text{NaHCO}_3$  buffer (pH 9.2) and ethanol. Endotoxin-free synthetic hemozoin, prepared as above, was added to the murine monocyte cell line RAW 264.7. The treated cells secreted a protein of

12.5 kDa which was shown to be MIF by Western blotting and N-terminal sequence analysis. Thus, hemozoin was a potent inducer of MIF in vitro at as little as 1 nmole heme equivalents. Furthermore, co-incubation of  
5 hemozoin with IFN- $\gamma$  at 10-100 U/ml further increased MIF release.

To further assess the role of MIF in malaria infection, blood was drawn from C3H/HeN mice exhibiting a low level of chronic parasitemia with  
10 *Plasmodium berghei* (10% parasitized red cells). Five  $\mu$ l serum aliquots were reacted with an anti-MIF antibody and analyzed by Western blotting. This relatively low level of parasitemia was associated with readily detectable levels of circulating serum  
15 MIF. Levels of parasitemia in the 10% range were sufficient to maintain persistent MIF production for as long as 28 days.

In addition, 500 nmoles of endotoxin-free synthetic hemozoin was injected i.p. into C3H/HeN mice  
20 and blood sampled 24 and 48 hours later. The sera were separated and assayed for MIF by Western blotting and TNF $\alpha$  by L929 cytotoxicity assays. Although circulating TNF $\alpha$  levels were undetectable in this study (<50 pg/ml), MIF was readily detected in sera at  
25 24 and 48 hours. Hence, a low level parasitemia was associated with circulating MIF in vivo. MIF appeared in sera under circumstances when TNF $\alpha$  was not detectable. Hemozoin alone was found sufficient to induce MIF production in vitro and in vivo. Taken  
30 together, these results show that MIF is an important early mediator in the host response to malaria infection. Furthermore, this observation indicates that anti-MIF therapy may be effective in ameliorating the pathological sequelae of parasite-induced cytokine  
35 release.

10. EXAMPLE: ANTI-MIF ABROGATES ANTIGEN-SPECIFIC IMMUNE RESPONSES

This example demonstrates that anti-MIF antibody can substantially reduce the magnitude of an antigen-specific immune response. This indicates that MIF is a critical component of an immune response, and thus anti-MIF treatment may be useful in substantially reducing an undesired immune reaction such as autoimmunity and allergy.

10.1. ANTI-MIF INHIBITS ANTIGEN-SPECIFIC T CELL PROLIFERATION

Ten BALB/c mice were primed in vivo with RNase A emulsified in complete Freund's adjuvant at 1:1 ratio. Each mouse received 20  $\mu$ g of RNase A in 0.2 ml i.p. After 2-3 weeks, splenic T cells were isolated and their proliferative activity assayed. In a microtiter plate,  $4 \times 10^5$  T cells were incubated with  $5 \times 10^5$  antigen-presenting cells per well. While the addition of RNase A led to an increased proliferation of the in vivo primed T cells as measured by  $^3\text{H}$ -thymidine incorporation, this proliferation was reduced to control levels when anti-MIF antibody was also added.

10.2. ANTI-MIF INHIBITS ANTIGEN-SPECIFIC INFLAMMATORY RESPONSE

BALB/c mice were primed in the left rear footpad and in the neck muscle with complete Freund's adjuvant at 0.2 ml per injection. After 10-14 days, the mice were boosted in the right rear footpad with 50  $\mu$ l of purified protein derivative (of tuberculin; PPD) with or without 50  $\mu$ l of anti-MIF antibody. When the width of the right rear footpads of the animals were measured with calipers 48 hours later as an indication of delayed-type hypersensitivity reaction, anti-MIF treatment was shown to reduce the swelling in the

footpads by greater than 50% as compared to controls. Analysis by light microscopy also revealed histopathological evidence of reduced inflammation in footpads of anti-MIF treated mice.

5

11. EXAMPLE: PRODUCTION OF MIGRATION INHIBITORY FACTOR BY THE PITUITARY GLAND

The following example describes the production of MIF by pituitary cells in vitro and in vivo.

10

11.1. MATERIALS AND METHODS

11.1.1. CELLS

The murine anterior pituitary cell line AtT-20 was obtained from American Type Culture Collection and maintained in low serum medium.

15

11.1.2. INDUCTION OF MIF

AtT-20 cells were cultured for 24 hours in serum-free medium (enriched serum-free Joklik's medium from Gibco) containing various concentrations of LPS (E. coli LPS 0111:B4 from Sigma) at 0.5  $\mu\text{g/ml}$  - 50  $\mu\text{g/ml}$ . Two ml of conditioned media was removed from  $1.5 \times 10^5$  cells and concentrated 100-fold by membrane filtration (10K cut off) prior to SDS polyacrylamide gel electrophoresis (18%). Silver staining analysis of the gels revealed a 12.5 kDa protein band. This protein was transferred to polyvinylidene difluoride membrane and its primary N-terminal amino acid sequence was determined by automated gas-phase sequencing (Applied Biosystems).

20

25

30

11.1.3. WESTERN BLOT

MIF release from pituitary (AtT-20) cells was analyzed by Western blotting with anti-MIF antibody (FIG. 8).  $1.5 \times 10^5$  pituitary cells were cultured

35



serum-free for 18 h with different concentrations of LPS. After transfer, nitrocellulose membranes were incubated with rabbit anti-rMIF serum (1:1000 dilution) and bound antibody visualized with  
5 horseradish peroxidase-conjugated goat anti-rabbit antibody. 100 ng of rMIF was electrophoresed on the same gel for reference. Control supernatants (C) contained 1  $\mu$ g/ml LPS that was added after removal of cells. Pre-immune serum showed no reactivity.

10

#### 11.1.4. IMMUNOCYTOCHEMISTRY

Cells were cultured for 16 h in the presence (FIG. 9B) or absence (FIG. 9A) of 25  $\mu$ g/ml LPS, washed, fixed with formaldehyde, and incubated with  
15 anti-rMIF serum (1:1000 dilution) prior to staining with immunoperoxidase-linked goat anti-rabbit antibody. No staining was observed with pre-immune serum or with immune serum that had been pre-incubated with rMIF.

20

#### 11.1.5. DETECTION OF MIF mRNA

Pituitary mRNA analyses from two representative mice are shown for each time interval (FIG. 10). RT-PCR analysis utilized murine MIF primers (22 cycles),  
25  $\beta$ -actin primers (25 cycles), or CD2 primers (pituitary cDNA: 35 cycles, control spleen cDNA: 25 cycles) as shown. Nine-week old mice were injected intraperitoneally with LPS at a dose of 2.25 mg/kg and sacrificed 0, 6, 16, 24 and 48 h later. Pituitary RNA  
30 was prepared by extraction with RNazol (Biotecx Laboratories, Inc.) from a total of four mice per time interval. After reverse transcription with oligo-dT, DNA amplification products were analyzed by agarose gel electrophoresis. Reverse transcription efficiency  
35 was assessed and normalized relative to  $\beta$ -actin mRNA.

The murine MIF primers were: 5'-CCATGCCTATGTTTCATCGTG-3' and 3'-GA-ACAGCGGTGCAGGTAAGTG-5' and were designed to amplify a 381 bp sequence that spanned at least one intron, as determined by Southern hybridization and internal restriction enzyme analysis of mouse genomic DNA. Murine intron-spanning CD2 primers were 5'-CCTGGTTCGCAGAGTTTAA-3' and 3'-TCTGTTCTTGCAAGACC-5', and  $\beta$ -actin primers were 5'-GTGGGCCGCTCTAGGCACCA-3' and 3'-TGGCCTTAGGGTTCAGGGGG-5'.

10

#### 11.1.6. POLYMERASE CHAIN REACTION

RNA was analyzed (FIG. 11) at 0 h and 24 h after LPS injection (2.25 mg/kg) by competitive PCR as described (Li et al., 1991, J. Exp. Med. 174:1259-1262), utilizing a 273 bp competitive MIF template. The competitive template was added in the range of 0.1 to 5 pg per reaction as shown in FIG. 11 and co-amplified with cDNA utilizing the MIF primers described in Section 9.1.5., supra. At 0 h, equivalent amounts of cDNA and competitor DNA amplification products appear in the lane showing 0.35 pg of MIF competitor template. At 24 h, equivalent amounts of cDNA and competitor DNA amplification products appear in the lane showing 1.0 pg of MIF competitor template.

25

#### 11.1.7. EXPRESSION OF MIF PROTEIN IN VIVO

Pituitary MIF protein content was measured in BALB/c and C3H/HeJ mice challenged with LPS (FIG. 12). A similar LPS-dependent decrease in pituitary MIF protein content was observed in both C3H/HeN and BALB/c<sup>nu/nu</sup> mice as in BALB/c mice. Mice (N = 2-5 per time point) were injected intraperitoneally with LPS at 2.25 mg/kg, pituitary lysates were prepared at intervals, and protein analyzed by Western blotting.

35

Aliquots representing  $\frac{1}{4}$  of a pituitary were electrophoresed, transferred to nitrocellulose membrane, and incubated with anti-rMIF serum (1:1000 dilution). Bound antibody was visualized by

- 5 incubation with horseradish peroxidase-conjugated goat anti-rabbit antibody and substrate.

- Nine-week-old mice were injected intraperitoneally with LPS at 2.25 mg/kg and serum samples (5  $\mu$ l each) obtained at indicated time
- 10 intervals were analyzed by Western blotting. Single bands corresponding to MIF were observed and quantified by laser densitometry (Shimadzu FDU-3, CS-9000 U). Recombinant MIF standards were electrophoresed and transferred in adjacent lanes.
- 15 Scanning integrals are presented as relative MIF content normalized to the respective 0 h zone on each membrane. Each plotted point is the mean  $\pm$  SEM obtained for individual sera from multiple animals (N = 2-5 per time point).

20

#### 11.2. RESULTS

The murine anterior pituitary cell line AtT-20 was cultured under low serum conditions in the presence of increasing amounts of bacterial endotoxin.

- 25 At time intervals, the conditioned media was collected, concentrated, and analyzed for the presence of secreted proteins by SDS polyacrylamide gel electrophoresis. Endotoxin stimulation was observed to result in the specific time- and
- 30 concentration-dependent secretion of a 12.5 kDa protein. The specificity of the secretory response was confirmed by the absence of LPS-induced cytopathicity, as determined by the lack of an LPS-related increase of supernatant lactate dehydrogenase
- 35 activity. These measurements revealed no

LPS-dependent increase in enzyme activity, despite prolonged (24 h) incubation under serum-free conditions (0  $\mu$ g LPS:  $112.8 \pm 17.9$  IU/L, 50 $\mu$ g LPS:  $119.7 \pm 13.1$  IU/L; P=NS). The 12.5 kDa protein was  
5 isolated, subjected to N-terminal microsequencing, and identified as the murine homolog of human MIF (96% identity over 27 amino acids).

Cellular activation by LPS is known to be potentiated by serum factors such as LPS-binding  
10 proteins that interact with specific cell surface receptors (Schuman et al., 1990, Science 349:1429-1431; Wright et al., 1990, Science 249:1431-1433). Supplementation of culture medium with 1% fetal bovine serum increased markedly the secretion of MIF from  
15 pituitary cells stimulated by LPS. As little as 100 pg/ml of LPS was found to induce pituitary cell secretion of MIF (FIG. 8). In contrast, MIF was not released by incubation with the inflammatory mediators tumor necrosis factor  $\alpha$ , (1-100 ng/ml), interleukin-1 $\beta$   
20 (1-100 ng/ml), interleukin-6 (1-100 ng/ml), or interferon- $\gamma$  (1-100 ng/ml). Western blotting analysis of pituitary cell lysates also revealed that resting, non-stimulated cells contained significant amounts of pre-formed MIF. Immunocytochemistry studies confirmed  
25 these results and showed the disappearance of substantial amounts of immunoreactive, intracellular MIF within 16 hr of LPS stimulation (FIG. 9A and 9B).

To investigate the pituitary expression of MIF in vivo, pituitary mRNA analysis was performed in mice  
30 injected with sublethal amounts of LPS (2.25 mg/kg). Pituitary RNA was isolated at increasing time intervals and subjected to reverse transcription-polymerase chain reaction (RT-PCR) analysis. Pituitary MIF mRNA levels increased with time and  
35 reached a plateau 16-24 h after LPS challenge in

endotoxin-sensitive mice (FIG. 10). These findings were confirmed by competitive PCR analysis of representative cDNA preparations that showed a 3-fold increase in pituitary MIF mRNA after LPS stimulation (FIG. 11). Although infiltrating mononuclear cells were not evident in stained pituitary sections obtained after LPS treatment, in order to exclude a potential T cell contribution of pituitary MIF mRNA, pituitary cDNA was amplified with primers specific for the T cell-specific gene product, CD2. Despite intentional overcycling, RT-PCR analysis of CD2 was uniformly negative, ruling out infiltrating T cells as a possible source of pituitary MIF mRNA (FIG. 10). LPS did not induce pituitary MIF mRNA in the genetically endotoxin-resistant strain C3H/HeJ, while congenic endotoxin-sensitive C3H/HeN mice showed an increase of pituitary MIF mRNA between 0 hr and 24 hr similar to that observed with BALB/c mice.

The pituitary content of MIF protein in vivo was analyzed by Western blotting of pituitary lysates. Pituitaries obtained from normal, non-stimulated mice showed significant amounts of pre-formed MIF protein (FIG. 12). A significant decrease in pituitary MIF content was observed 8-20 h after LPS injection in endotoxin-sensitive mice, but not in endotoxin-resistant (C3H/HeJ) mice.

Previous studies over the years have identified MIF to be a product of lectin-stimulated T cells (Weiser et al., 1989, Proc. Natl. Acad. Sci. USA 86:7522-7526; Weiser et al., 1981, J. Immunol. 126:1958-1962). Since LPS does not directly activate T cells, MIF that appears in the circulation presumably results from direct stimulation of various cell types, including but not limited to pituitary cells and monocyte/macrophages by LPS or from indirect

activation LPS-induced mediators of these and other cell types, including for instance T cells. In order to better establish the contribution of various cell types to circulating serum MIF in endotoxemic mice, control BALB/c, and T cell-deficient BALB/c<sup>nu/nu</sup> mice were injected intraperitoneally with LPS, serum samples were analyzed by gel electrophoresis and Western blotting, and serum MIF levels quantified by laser densitometry (FIG. 13). Serum MIF was detected at 2 h in wild-type mice (BALB/c) and increased over 20 h in a time-dependent fashion. In contrast, T cell-deficient (nude) mice showed a markedly delayed response in the appearance of serum MIF. MIF levels increased significantly only after 8 h of LPS challenge.

These findings indicate that pituitary MIF contributes directly to circulating serum MIF that is detectable in the post-acute phase ( $\geq 8$  hours) of endotoxaemia. This time course is consistent with the disappearance of pituitary MIF protein and the induction of pituitary MIF mRNA levels that were observed in vivo (FIGS. 10 and 11). The results obtained in nude mice indicate that T cell MIF contributes primarily to the serum MIF that appears in the first 8 h. LPS treatment (2.25 mg/kg) of endotoxin-resistant C3H/HeJ mice produced no detectable serum MIF.

## 12. EXAMPLE: PRODUCTION OF MIGRATION INHIBITORY FACTOR BY MACROPHAGES

### 12.1. MATERIALS AND METHODS

#### 12.1.1. REAGENTS

*E. coli* O111:B4 LPS, polymyxin B, carbenicillin, PMSF and Tween-20 were obtained from Sigma (St. Louis,

MO). LPS was resuspended in pyrogen-free water, vortexed, sonicated, aliquoted (5 mg/ml), and stored at -20°C. Serial dilutions of LPS were prepared in pyrogen-free water by sonication (Branson 3210, Danbury, CT). Gentamicin was from Gibco (Grand Island, NY). Thioglycollate broth (Difco, Detroit, MI) was prepared according to the manufacturer's recommendation, autoclaved, and stored protected from light at room temperature. Horseradish peroxidase-conjugated goat anti-rabbit antibody was purchased from Pierce (Rockford, IL) and 4-chloro-1-naphthol and stabilized 3,3',5,5'-tetramethylbenzidine (TMB) substrate for horseradish peroxidase were from Promega (Madison, WI). Polyclonal anti-MIF serum was generated by immunizing New Zealand White rabbits (Hare Marland, Hewitt, NJ) with purified recombinant murine MIF. On week 1 and 2, rabbits were inoculated intra-dermally with 100 µg of rMIF diluted in complete Freund's adjuvant, and with 50 µg of rMIF diluted in incomplete Freund's adjuvant on week 4. Immune serum was collected one week after the last inoculation.

#### 12.1.2. CYTOKINES

Recombinant murine MIF (rMIF) was expressed in *E. coli* BL21/DE3 (Novagen, Madison, WI) and purified to homogeneity by anion exchange (Mono Q; Pharmacia, Piscataway, NJ) and reverse phase chromatography (C8-SepPak, Millipore, Milford, MA), lyophilized and reconstituted in sodium phosphate buffer (20 mM, pH 7.2) following procedures in Example 6, supra. MIF bioactivity was established by measuring dose-dependent MIF-induced augmentation of *Leishmania major* killing by macrophages and by neutralization of this activity with anti-MIF antibody. rMIF contained 9 pg of endotoxin per µg protein as determined by the

chromogenic *Limulus* amoebocyte assay (Bio-Whittaker Inc., Walkersville, MD). Recombinant murine IL-1 $\beta$  and IL-6 (5  $\mu$ g/ml after reconstitution) were obtained from R&D (Minneapolis, MN) and recombinant murine IFN $\gamma$  (10<sup>5</sup> IU/ml) was from Boehringer-Mannheim (Indianapolis, IN). Cytokines were reconstituted in pyrogen-free water containing 0.1% of very low endotoxin BSA (Miles Inc., Kankakee, IL) and stored at -80°C. The endotoxin content of the reconstituted cytokines was 0.5 ng/ $\mu$ g of TNF $\alpha$ , 2 ng/ $\mu$ g of IL-1 $\beta$ , 1.1 ng/ $\mu$ g of IL-6 and 0.06 pg/unit of IFN $\gamma$  as determined by the chromogenic *Limulus* amoebocyte assay.

#### 12.1.3. ANIMALS AND EXPERIMENTAL DESIGN

Eight to 10-week-old (19-21 g) female BALB/c (control), T cell-deficient BALB/c<sup>nu/nu</sup> (nude), and hypophysectomized BALB/c mice were purchased from Charles River (Kingston, NY). Animals were housed in groups of 5 or 10 mice per cage with free access to food and water (supplemented with 5% glucose for hypophysectomized mice) and were acclimatized for 5 days prior to experimentation. Mice were injected i.p. with nonlethal doses of LPS (2.25 mg/kg for BALB/c and T cell-deficient mice, and 50  $\mu$ g/kg for hypophysectomized mice), and sacrificed 2, 8 and 20 h after LPS challenge to collect serum. The LPS dose was adjusted to provide a comparable degree of lethality in the control BALB/c group as among the hypophysectomized mice, which are hypersensitive to endotoxin. Five  $\mu$ l of serum was analyzed by Western blotting and visualized with anti-MIF antibody.



12.1.4. CELLS

RAW 264.7 murine macrophages, THP-1 human monocytes, ASL-1 murine and Jurkat human T lymphocyte cell lines were obtained from the American Type Culture Collection (Rockville, MD). Cells were grown in RPMI 1640 medium (Gibco, Grand Island, NY) containing 2 mM glutamine, 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT), 50 µg/ml of carbenicillin and gentamicin. Medium for ASL-1 cells additionally was supplemented with 1 mM sodium pyruvate. Thioglycollate-elicited peritoneal macrophages were obtained from BALB/c mice that were injected i.p. 3-4 days previously with 2 ml of sterile thioglycollate broth. Cells were harvested under strict aseptic conditions by lavage of the peritoneal cavity with 5 ml of an ice-chilled 11.6% sucrose solution. After centrifugation (10 min at 800 g), cells were resuspended in RPMI/10% FBS, enumerated, and plated at a density of  $2 \times 10^6$  or  $4 \times 10^6$  cells/well. Human polymorphonuclear leukocytes were isolated from peripheral blood by density gradient centrifugation (Vadas et al. 1979, J. Immunol. 122: 1228; Sherry et al., 1981, In: Methods for Studying Mononuclear Phagocytes, Academic Press, NY, p. 187).

25

12.1.5. MIF CONTENT OF MACROPHAGE  
AND T LYMPHOCYTE LYSATES

Aliquots of  $1 \times 10^6$  cells of each type were lysed with Tris-buffered saline (50 mM Tris-Base, 150 mM NaCl, pH 7.5) containing 1% NP-40, 0.5% deoxycholic acid, 0.1% SDS and 2 mM EDTA. Cellular debris was pelleted and the supernatants were diluted with an equal volume of reducing SDS-PAGE sample buffer. Ten microliter of lysate (equivalent to  $5 \times 10^3$  cells) were electrophoresed through 18% polyacrylamide gels, and

35

transferred to nitrocellulose membranes for Western blot analysis using polyclonal rabbit anti-MIF serum.

12.1.6. STIMULATION OF MACROPHAGES  
BY LPS AND CYTOKINES

5 RAW 264.7 macrophages were washed with fresh medium, harvested by gentle scraping, resuspended in RPMI/10% FBS, and incubated at  $2 \times 10^6$  or  $4 \times 10^6$  cells/well in 3.5-cm tissue culture plates (Linbro®,  
10 Flow, McLean, VA). After 3 h of incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, nonadherent cells were removed and wells were washed twice with RPMI/1% FBS. Cells then were incubated for 12 h with LPS (at concentrations ranging from 1 pg/ml to 1 µg/ml) or  
15 with cytokines (1 or 10 ng/ml for TNFα, IL-1β, and IL-6, and 10, 100 or 1000 IU/ml for IFNγ) diluted in RPMI/1% FBS. For time-course experiments, conditioned media of parallel cultures were removed at 3, 6, 9, 12 and 15 h intervals after LPS stimulation.  
20 Thioglycollate-elicited peritoneal macrophages ( $10^7$  cells/well) were harvested as described and conditioned similarly to RAW 264.7 cells. Cells were incubated for 12 h either with LPS or with IFNγ plus LPS. When co-stimulated with IFNγ and LPS, cells were  
25 first incubated for an hour with IFNγ (100 IU/ml) before addition of LPS at the indicated concentrations. At the end of each experiment, cell-culture media were collected, centrifuged (10 min at 800 g), and supplemented with PMSF (1 mM).  
30 Supernatants then were concentrated 10-fold by membrane filtration (10 kD cut-off) (Centricon-10, Amicon, Beverly, NY). Cellular RNA was extracted from adherent cells.

35

12.1.7. REVERSE TRANSCRIPTION AND  
POLYMERASE CHAIN REACTION  
(RT-PCR)

Total cellular RNA was extracted from macrophages with RNazol-B (Tel-Test Inc., Friendswood, TX) following the manufacturer's protocol. One  $\mu$ g of RNA was reversed transcribed using oligo (dT)<sub>12-18</sub> and M-MLV reverse transcriptase (Gibco, Grand Island, NY) in a 50  $\mu$ l reaction. Five  $\mu$ l of cDNA was amplified by PCR in a Perkin-Elmer/Cetus 9600 thermal cycler (denaturation 1 min at 94°C, annealing 1 min at 55°C, elongation 1 min at 72°C) using murine MIF primers (32 cycles), TNF $\alpha$  primers (22 cycles), or  $\beta$ -actin primers (25 cycles). Intron-spanning TNF $\alpha$  primers were 5'-GCGGAGTCCGGGCAGGTCTA-3' and 3'-GGGGGGCTGGCTCTGTGAGG-5'. DNA amplification products were analyzed on 2% agarose gels and gel loading was normalized to  $\beta$ -actin PCR products. Quantification of MIF mRNA from LPS-stimulated RAW 264.7 macrophages was done by competitive PCR (Gilliland et al., 1990, Proc. Natl. Acad. Sci. USA 87: 2275). The 273 bp competitive cDNA template was added at the indicated concentrations and co-amplified with MIF cDNA using MIF primers.

12.1.8. WESTERN BLOTS

For Western blotting, samples were resolved on 18% SDS polyacrylamide gels under reducing conditions and transferred onto nitrocellulose membrane (Schleicher and Schuell, Keene, NH) at 50 V and 150 mA for 16 h. Membranes were blocked with 500 mM NaCl, 50 mM Tris-Base, pH 7.5 buffer containing 5% nonfat dry milk and 0.05% Tween-20 and incubated first with polyclonal rabbit anti-rMIF serum and then with horseradish peroxidase-conjugated goat anti-rabbit IgG antibody (each diluted 1:1000 in 500 mM NaCl, 50 mM Tris-Base, pH 7.5 buffer with 1% BSA and 0.05%

Tween-20). Polyclonal rabbit anti-murine MIF serum was demonstrated to react with both natural and recombinant murine and human MIF. Incubations were for 1 h each. MIF was visualized by development with  
5 chloronaphthol/H<sub>2</sub>O<sub>2</sub> or with TMB substrate. Serum MIF bands were quantified by laser densitometry (Shimadzu FDU-3, CS-9000U, Braintree, MA). Faint densitometric signals in the MIF region were detected in most sera prior to LPS administration (0 h). Hence, scanning  
10 integrals are presented as relative serum MIF content (i.e. fold-increase) normalized to the respective 0 h zone (i.e. background or baseline staining) on each membrane. Each plotted point is the mean  $\pm$  SEM of individual sera from 2 to 5 animals.

15

#### 12.1.9. MIF-INDUCED SECRETION OF TNF $\alpha$ BY MACROPHAGES

Cells (4x10<sup>6</sup>/well) were processed as described above and stimulated for 12 h with rMIF at specified concentrations ranging from 100 pg/ml to 1  $\mu$ g/ml. At  
20 1  $\mu$ g/ml of rMIF, LPS contamination was 9 pg/ml as assessed by *Limulus* assay. To neutralize this minute amount of LPS, rMIF was preincubated for 1 h at 37°C with polymyxin B (1  $\mu$ g/ml). MIF-induced TNF $\alpha$  activity in cell-culture supernatant was quantitated by L929  
25 cell cytotoxicity (Wolpe et al., 1988, J. Exp. Med. 167: 570). MIF-induced TNF $\alpha$  activity was blocked completely by anti-TNF $\alpha$  antibodies. rMIF did not contribute to TNF $\alpha$  activity, as recombinant murine TNF $\alpha$  (dose range: 5 pg/ml to 1  $\mu$ g/ml) cytotoxicity  
30 remained unchanged when rMIF (10 pg/ml to 10  $\mu$ g/ml) or anti-MIF polyclonal serum were added to L929 cells together with rTNF $\alpha$ .

35

## 12.2. RESULTS

### 12.2.1. SERUM MIF LEVELS IN LPS-INJECTED MICE

Pituitary MIF contributes significantly to the MIF that appears in serum during endotoxemia. Serum MIF reached peak levels in normal mice at 20 h but was undetectable in hypophysectomized mice at this time, demonstrating that the pituitary is a major source of the MIF that appears in serum after LPS injection.

The central role played by MIF in endotoxic shock, together with the known hypersensitivity of hypophysectomized mice to LPS, led to experiments to examine more closely the kinetics of MIF appearance in the circulation during the early, acute phase (i.e. 2-8 h) of endotoxemic shock. For comparison purposes, serum MIF kinetics were also examined in nude mice, which lack a T cell source of MIF. Control, hypophysectomized, and nude mice were injected with LPS and serum prepared at intervals and quantified for MIF content by Western blotting and laser scanning densitometry (FIG. 14). Whereas serum MIF concentrations increased gradually over 20 h in control mice, hypophysectomized mice exhibited a rapid rise and then prompt fall in MIF that peaked 2 h after LPS injection. In nude mice, the rise in serum MIF was delayed by 8 to 12 h, but increased in a manner that was similar to that of the control mice. Given the lack of a pituitary source of MIF in the hypophysectomized mice, the MIF present in the circulation at 2 h reflected the release of MIF by an additional LPS-sensitive cell population. Such serum MIF kinetics were reminiscent of a macrophage TNF $\alpha$  response (Beutler et al., 1985, J. Exp. Med. 161: 984; Beutler et al., 1985, J. Immunol. 135: 3972; Michie et al., 1988, N. Eng. J. Med. 318: 1481) and thus the

possibility that MIF was produced by cells of the monocyte/macrophage lineage was examined.

#### 12.2.2. EXPRESSION OF MIF IN RESTING MACROPHAGES

5        Since both a pituitary cell-line (AtT-20) and the whole pituitary *in vivo* contain substantial amounts of pre-formed MIF, the intracellular MIF content of various inflammatory cells and cell lines was analyzed. Significant amounts of pre-formed MIF  
10 protein were found to be present in cell lysates obtained from resting, non-stimulated murine RAW 264.7 cells, murine peritoneal macrophages, and human THP-1 monocytes (FIG. 15). It was estimated that on average there was 0.1-1 pg of immunoreactive, pre-formed MIF  
15 protein per macrophage. The MIF content of murine or human monocytes/macrophages was similar to that of two T cell lines, the human Jurkat and the murine ASL-1. In contrast, cell lysates obtained from purified polymorphonuclear leukocytes (PMNs) exhibited no  
20 detectable MIF protein (FIG. 15).

#### 12.2.3. MIF SECRETION BY LPS-STIMULATED MACROPHAGES

MIF was observed to be secreted by macrophages into culture medium after LPS stimulation. In RAW  
25 264.7 macrophages, MIF secretion was induced by as little as 10 pg/ml of LPS, peaked at 1 ng/ml, and was not detectable at LPS concentrations > 1 µg/ml (FIG. 16A). In elicited peritoneal macrophages, 1 ng/ml of LPS was required to induce MIF secretion, and a  
30 maximal response was observed with 10-100 ng/ml of LPS (FIG. 16B). Of significance, co-stimulation of cells with IFNγ (100 IU/ml, given 1 h prior to LPS) plus LPS resulted in a > 1000-fold increase in LPS  
responsiveness (FIG. 16C).

35

We next examined the time-course of MIF secretion in parallel cultures of RAW 264.7 macrophages maximally stimulated with 1 ng/ml of LPS. As assessed by Western blotting, MIF was first detected 6 to 9 h after LPS stimulation. Maximum amounts of MIF appeared in medium between 9 and 12 h post-LPS stimulation. MIF levels then decreased, indicating that by 12 h, MIF is removed or degraded at a rate that exceeds that of synthesis and release by macrophages.

#### 12.2.4. EXPRESSION OF MIF mRNA BY LPS-STIMULATED MACROPHAGES

The expression of MIF mRNA by LPS-stimulated RAW 264.7 macrophages was investigated by reverse transcription and polymerase chain reaction (RT-PCR). Parallel cultures were incubated for 12 h with medium or increasing amounts of LPS and analyzed for the expression of MIF, TNF $\alpha$ , and  $\beta$ -actin. MIF mRNA was expressed constitutively in non-stimulated murine RAW 264.7 cells (FIG. 17) and in elicited peritoneal macrophages. As expected, the expression of TNF $\alpha$  mRNA increased over the range of LPS concentrations (1 pg/ml to 1  $\mu$ g/ml). In contrast, MIF mRNA levels correlated inversely with LPS concentration over this dose range. MIF mRNA level was highest in cells induced with 1 pg/ml of LPS and lowest in those induced by 1  $\mu$ g/ml of LPS.

To assess more quantitatively the induction of MIF mRNA, cDNA was prepared from control and LPS-stimulated cultures of RAW 264.7 macrophages and analyzed by competitive PCR. The amount of competitive template which was required to obtain equivalent levels of MIF and competitor DNA amplification products was 1.5 pg for non-stimulated (control) macrophages and 3 pg for LPS-induced

(100 pg/ml) RAW 264.7 macrophages (FIG. 18). Therefore, macrophage MIF mRNA increased approximately 2-fold after LPS induction, an increase comparable to that observed in pituitary cells *in vivo* (3-fold after  
5 induction with 45  $\mu$ g of LPS injected i.p.). The time-course of MIF mRNA induction was examined in RAW 264.7 macrophages incubated with 1 ng/ml of LPS, the LPS concentration which induced maximum MIF secretion in these cells. MIF mRNA levels increased 6 h after LPS  
10 induction and remained elevated for up to 12 h post-LPS.

#### 12.2.5. INTERACTION BETWEEN MIF AND PRO-INFLAMMATORY CYTOKINES

15 The detection of elevated levels of MIF in blood after LPS administration together with the finding of an important role for MIF in experimental endotoxemia led to the investigation of the interaction between MIF and other pro-inflammatory cytokines. MIF release  
20 by RAW 264.7 macrophages was studied after stimulation with recombinant murine TNF $\alpha$ , IL-1 $\beta$ , IL-6, or IFN $\gamma$ . By Western blotting, TNF $\alpha$  and IFN $\gamma$  were found to induce MIF secretion in a dose-dependent fashion (FIG. 19). The lowest concentration of cytokine which was  
25 effective under these experimental conditions was 1 ng/ml of TNF $\alpha$  and 10 IU/ml of IFN $\gamma$ . The TNF $\alpha$  and IFN $\gamma$  effects could not be accounted for by LPS contamination of recombinant cytokine preparations, which were found to be 0.5 pg/ng of TNF $\alpha$  and 60 fg/U  
30 of IFN $\gamma$ . LPS and IFN $\gamma$  had additive/synergistic effect on MIF secretion, whereas TNF $\alpha$  and IL-1 $\beta$  did not. Finally, MIF production was not induced by either IL-1 $\beta$  or IL-6 at 1 or 10 ng/ml.

The secretion of TNF $\alpha$  and IL-1 $\beta$  by RAW 264.7  
35 macrophages was examined after stimulation with rMIF. MIF samples were pre-incubated for 1 h with



polymyxin B at 1  $\mu\text{g/ml}$  to neutralize small amounts of contaminating LPS (9  $\text{pg}/\mu\text{g}$  of MIF by the chromogenic *Limulus* assay). rMIF at concentrations  $\geq 100 \text{ ng/ml}$  was found to induce the secretion of bioactive  $\text{TNF}\alpha$  as determined by the L929 cytotoxicity assay (FIG. 20). The MIF-induced  $\text{TNF}\alpha$  bioactivity was blocked completely by anti- $\text{TNF}\alpha$  antibodies and rMIF did not contribute to  $\text{TNF}\alpha$  cytotoxicity, as recombinant murine  $\text{TNF}\alpha$  cytotoxicity was unchanged when rMIF or anti-MIF polyclonal serum were added to L929 cells together with r $\text{TNF}\alpha$ . IL- $1\beta$  secretion, in contrast, was not detectable by Western blotting over the concentration of rMIF tested.

**15**           **13.**     **EXAMPLE:**    **STEROID INDUCES MIF**  
                      SECRETION BY MACROPHAGES

### 13.1. MATERIALS AND METHODS

20 13.1.1. CELL PREPARATION AND MIF INDUCTION

RAW 264.7 murine macrophages were obtained from the American Type Culture Collection (Rockville, MD). Cells were grown in RPMI 1640 medium (Gibco, Grand Island, NY), 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT), 50  $\mu$ g/ml of carbenicillin and gentamicin. RAW 264.7 macrophages were washed with fresh medium, harvested by gentle scraping, resuspended in RPMI/10%FBS, and incubated at  $3 \times 10^6$  cells/well in 3.5-cm tissue culture plates (Linbro\*, Flow, McLean, VA). After 3h of incubation at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>, nonadherent cells were removed and wells were washed twice with RPMI/1%FBS. In some experiments designed to test for inhibition of MIF release, test compounds were then added to the culture media. For example, the cells were pre-treated with 20 $\alpha$ -dihydrocortisol at a final

concentration of  $10^{-6}M$ . Cells then were incubated for 12-15h with various steroids at concentrations ranging from  $10^{-6}M$  to  $10^{-14}M$  (e.g., dexamethasone), along with the desired test compound (e.g.,  $20\alpha$ -dihydrocortisol) if any. At the end of experiment, cell culture media were collected and centrifuged (10 min at 800 g). Cell-conditioned media were concentrated 10-fold by membrane filtration (10 kDa cut-off) (Centricon-10, Amicon, Beverly, NY).

- 10 Thioglycollate-elicited peritoneal macrophages were obtained from BALB/c mice that were injected i.p. 3-4 days previously with 2 ml of sterile thioglycollate broth. Cells were harvested under strict aseptic conditions by lavage of the peritoneal cavity with 5 ml of an ice-chilled 11.6% sucrose solution. After centrifugation (10 min at 800 g), cells were resuspended in RPMI/10% FBS, enumerated and plated at a density of  $3-4 \times 10^6$  cells/well. Cells were incubated for 12 to 15h with various steroids (at concentrations ranging from  $10^{-6}M$  to  $10^{-14}M$ ).

- Samples were resolved on 18% SDS polyacrylamide gels under reducing conditions and transferred onto nitrocellulose membrane (Schleicher and Schuell, Keene, NH) at 50 V and 150 mA for 16 h. Membranes were blocked with 500 mM NaCl, 50 mM Tris-Base, pH 7.5 buffer containing 5% nonfat dry milk and 0.05% Tween-20 and incubated first with polyclonal anti-rMIF serum and then with horseradish peroxidase-conjugated goat anti-rabbit IgG antibody (each diluted 1:1000 in 500 mM NaCl, 50 mM Tris-Base, pH 7.5 buffer with 1% BSA). Polyclonal rabbit anti-murine MIF was shown to react with both natural and recombinant murine and human MIF. Incubations were 1 h each. MIF was visualized by development with chloronaphtol/ $H_2O_2$  or with TMB substrate.

13.1.2. STEROID INJECTION

Canulated Sprague-Dawley female rats were purchased from Zivic-Miller. Animals were housed with free access to food and water and were acclimatized for at least 2 days prior to experimentation. Rats were injected i.v. through the jugular vein with either dexamethasone (1, 10 or 20 mg/kg), or with RU38486 (mifepristone) (1 or 10 mg/kg), or with the two drugs together (both at 10 mg/kg, RU38486 was injected 15 min. before dexamethasone). Serum was collected 3, 6, and 9 hours post-steroids injection and analyzed by Western blotting with anti-MIF antibody.

15

13.2. RESULTS

MIF was observed to be secreted by murine macrophages (RAW 264.7 macrophages and/or elicited peritoneal macrophages) after stimulation with dexamethasone, hydrocortisol, 5 $\beta$ -Pregnane-3 $\alpha$ ,11 $\beta$ ,17 $\alpha$ ,21-tetrol-20-one (tetrahydrocortisol), 4-pregnene-11 $\beta$ ,17 $\alpha$ ,20 $\alpha$ ,21-tetrol-3-one (20 $\alpha$ -dihydrocortisol), RU38486 (mifepristone), aldosterone, testosterone, and progesterone (Table V). In addition, dexamethasone at 10 mg/kg, RU 38486 at 10mg/kg or a combination of both drugs induced MIF secretion in vivo. Serum MIF was detected 3 to 6 hours after steroid injection.

As shown in FIG. 21, pretreatment and co-administration of 20 $\alpha$ -dihydrocortisol attenuates or abolishes the MIF response to dexamethasone.

35

TABLE V. STEROIDS INDUCE MIF SECRETION BY MURINE MACROPHAGES

Steroids	Range of Steroid Concentrations (M) Inducing MIF in:	
	RAW 264.7 Macrophages	Peritoneal Macrophages
Dexamethasone	$10^{-8}$ to $10^{-14}$	$10^{-8}$ to $10^{-12}$
Hydrocortisone	$10^{-8}$ to $10^{-12}$	$10^{-8}$ to $10^{-16}$
5 $\beta$ -Pregnane -3 $\alpha$ ,11 $\beta$ ,17 $\alpha$ ,21- Tetrol-20-one (Tetrahydrocortisol)	$10^{-8}$ to $10^{-14}$	Not tested
4-pregnene -11 $\beta$ ,17 $\alpha$ ,20 $\alpha$ ,21-tetrol-3-one (20 $\alpha$ dihydrocortisol)*	$10^{-10}$ to $10^{-14}$	$10^{-10}$ to $10^{-14}$
Aldosterone	$10^{-10}$ to $10^{-14}$	$10^{-8}$ to $10^{-14}$
Testosterone	$10^{-6}$ to $10^{-14}$	$10^{-8}$ to $10^{-14}$
RU-38486 (mifepristone)*	$10^{-8}$ to $10^{-14}$	Not tested
Progesterone	$10^{-4}$ to $10^{-12}$	Not tested

\* At higher concentrations, these steroids inhibit the macrophage release of MIF in response to a challenge dose of dexamethasone, i.e., a dose that normally would induce MIF release.

14. EXAMPLE: ORGAN DISTRIBUTION OF MIF  
AND MIF RECEPTOR

The following example describes the organ distribution of MIF and MIF receptor in mice.

5

14.1. MATERIALS AND METHODS

14.1.1. LABELING OF EXOGENOUS MIF

Five  $\mu\text{l}$   $^{125}\text{I}$ -MIF (specific activity,  $2 \times 10^4$  cpm/ng) together with 5  $\mu\text{l}$   $^{51}\text{Cr}$ -labeled red blood cells (RBCs), were injected intravenously in a total of 300  $\mu\text{l}$  saline. One preparation additionally contained 40  $\mu\text{g}$  unlabeled MIF as a competitor. Ten minutes prior to injection, each mouse was given 10 mg Na-iodide i.p. Mice were sacrificed 10 minutes after injection.

15

14.1.2. TISSUE PREPARATION

Four hundred  $\mu\text{l}$  blood was collected from each mouse immediately after sacrificing. Additionally, various organs were removed, rinsed in PBS, and blotted dry on filter paper. Organs were weighed and radioactivity was counted on two separate windows (one for  $^{125}\text{I}$ , one for  $^{51}\text{Cr}$ ) on a gamma counter. The remainder of the carcass was also weighed and counted.

25

14.1.3. ORGAN DISTRIBUTION OF ENDOGENOUS MIF

The organ distribution of MIF protein was examined in a 8 to 10-week old female BALB/c mouse sacrificed by  $\text{CO}_2$  asphyxiation and necropsied under aseptic conditions. Organs (brain, liver, spleen, kidneys, adrenals and lungs) were excised, sectioned and homogenized at  $4^\circ\text{C}$  with Tris-buffered saline (50 mM Tris-Base, 150 mM NaCl, pH 7.5) containing 1% NP-40, 0.5% deoxycholic acid, 0.1% SDS, 2 mM EDTA and 1 mM PMSF. Cellular debris was pelleted and aliquots of organ lysate supernatants, adjusted for protein

35

concentration, were diluted with an equal volume of reducing SDS-PAGE sample buffer. Samples (equivalent to 60  $\mu$ g of protein) were electrophoresed through 18% polyacrylamide gels and transferred to nitrocellulose  
5 membranes for Western blot analysis using polyclonal anti-MIF serum.

#### 14.2. RESULTS

The organ distribution of endogenous MIF protein  
10 was examined in normal mice. Mouse tissues were homogenized and aliquots of total protein electrophoresed and analyzed by Western blotting. As shown in FIG. 22, pre-formed MIF was detectable in mouse liver, spleen, kidney and brain. Much lower  
15 amounts were also found in the adrenals and lungs. These organs contain significant amounts of macrophages, suggesting that the MIF protein present in these tissues is at least partly macrophage-associated.

20  $^{125}$ I-MIF was also injected into mice in order to determine the tissue distribution of MIF receptors. The index of the tissue specific localization of radiolabelled MIF was determined, as described in Beutler et al., 1985, J. Immunol. 135:3972. As shown  
25 in Table VI, a particularly high level of MIF-binding was observed in the kidneys and liver among the organs examined.

30

35

TABLE VI

5	Organs	Index of Tissue Specific Localization of Radiolabelled MIF
	Liver	256
	Kidney	2334
	Lung	52
10	Adrenal	47
	Spleen	78
	Small Intestine	247
	Large Intestine	122
	Brain	17
15	Skin	103

#### 15. EXAMPLE: MIF RECEPTOR IDENTIFICATION

In this example, receptor proteins that bind to  
20 MIF are identified, and the partial amino acid  
sequences of each of these MIF receptors are  
presented.

##### 15.1. MATERIALS AND METHODS

25

##### 15.1.1. CELL LYSATE PREPARATION

Two large tissue culture plates ( $5 \times 10^7$  cells) of  
murine RAW 264.7 cells (a monocyte-like line) were  
solubilized by the addition of 2.5 ml of lysis buffer  
30 (50 mM Tris pH 7.5, 150 mM NaCl, 1% Triton X-100, with  
protease inhibitors). Plates were incubated on ice  
with gentle rotation. Plates were then scraped, the  
contents collected, and centrifuged in 4 microfuge  
(1.5 ml) tubes at 10,000 x g for 10 min at 4°C.  
35 Supernatants were collected and stored on ice.

#### 15.1.2. AFFINITY CHROMATOGRAPHY

For preparation of an MIF-conjugated affinity column, 1 gm of CNBr-Sepharose was first washed with 1 mM HCl. Bacterial recombinant murine MIF (3 mg) was  
5 dissolved in 40% acetonitrile/H<sub>2</sub>O. MIF solution then was mixed with activated CNBr-Sepharose and incubated for 3 hr at room temperature. Unreacted CNBr-Sepharose then was quenched by incubation with 0.2 M glycine pH 8.0 overnight. Coupling efficiency was  
10 determined by comparison of optical absorbance of MIF solution before and after coupling, and determined to be 85%. Resin then was washed, poured into a column and equilibrated with lysis buffer.

Supernatants of lysed RAW cell preparation were  
15 loaded onto MIF-Sepharose at a flow rate of 2 ml/hr, and washed sequentially with 3 column volumes of lysis buffer. Remaining bound protein was eluted with 5 volumes of 0.05% Triton X-100 in 0.1 M glycine, pH 2.5. Fractions were collected and buffered with 90 µl  
20 of 1 M Tris base.

#### 15.1.3. GEL ELECTROPHORESIS

Fractions containing protein species of interest were concentrated over Centricon-10 membranes,  
25 resuspended in equal volumes of 2X Laemmli sample buffer, and electrophoresed on 7.5% SDS-PAGE gels. Proteins were transferred onto PVDF filters utilizing 10 mM CAPS pH 11/20% methanol transfer buffer. Filters were stained with Ponceau-S and protein bands  
30 corresponding to molecular weights 52 kD and 72 kD were identified and excised.



#### 15.1.4. MICROSEQUENCING ANALYSIS

The excised protein bands were subjected to microsequencing analysis using automated gas-phase sequencing methods (Applied Biosystems).

5

#### 15.2. RESULTS

Putative MIF receptor protein was isolated by MIF affinity chromatography of RAW cell lysates on an MIF affinity column, as described above in Section 13.1.2.

10 Three fractions (6, 7, 8) were found to contain two protein species, 52 kD and 72 kD, when analyzed by SDS-PAGE. When parallel transfers were performed onto membranes which were then incubated with <sup>125</sup>I-labelled MIF, it was demonstrated that both the 52 kD and the  
15 72 kD bands bound radiolabelled MIF. Ligand binding was inhibited by simultaneous incubation of membranes with excess unlabelled MIF, thereby showing that binding was MIF specific.

Protein from the 52 kD species was isolated and  
20 subjected to amino acid microsequencing. A partial amino acid sequence was obtained from this 52 kD putative MIF receptor, as shown here:  
Ile-X-His-Asn-Thr-Val-Ala-Thr-Glu-Ile-(Ser)-(Gly)-Tyr-Asn-(Asn/Gly)-(Ala)-(Met)

25 The residues in parenthesis are tentative designation.

Similarly, protein from the 72 kD species was isolated and sequenced. A partial amino acid sequence was determined and shown here:

30 Ala-Lys-Lys-Gly-Ala-Val-Gly-Gly-Ile

The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the  
35 invention. Indeed, various modifications of the

invention in addition to those shown and described  
herein will become apparent to those skilled in the  
art from the foregoing description and accompanying  
drawings. Such modifications are intended to fall  
5 within the scope of the appended claims.

All publications cited herein are incorporated by  
reference in their entirety.

10

15

20

25

30

35

**WHAT IS CLAIMED IS:**

1. A method for treating a condition involving cytokine-mediated toxicity comprising administering to  
5 an individual an effective amount of an agent capable of inhibiting MIF biologic activity.
2. The method of Claim 1 in which the agent is an MIF antagonist.  
10
3. The method of Claim 2 in which the antagonist is an anti-MIF antibody.
4. The method of Claim 2 in which the  
15 antagonist is a soluble MIF-receptor.
5. The method of Claim 2 in which the antagonist is a small organic molecule.
- 20 6. A method for treating a condition involving cytokine-mediated toxicity comprising administering to an individual an effective amount of an agent capable of inhibiting MIF receptor biologic activity.
- 25 7. The method of Claim 6 in which the agent is an MIF-receptor antagonist.
8. The method of Claim 7 in which the antagonist is an anti-MIF-receptor antibody.  
30
9. The method of Claim 7 in which the antagonist is a biologically inactive MIF analog which is capable of binding to MIF receptor.  
35

10. The method of Claim 7 in which the antagonist is a small organic molecule.

11. A method for treating a condition involving  
5 cytokine-mediated toxicity comprising administering to an individual an effective amount of an agent capable of inhibiting MIF gene expression.

12. The method of Claim 11 in which the agent is  
10 an antisense molecule complementary to MIF mRNA.

13. The method of Claim 11 in which the agent is a ribozyme molecule specific for MIF mRNA.

14. The method of Claim 11 in which the agent is  
15 a triple helix component.

15. The method of Claim 11 in which the agent is a steroid.  
20

16. A method for treating a condition involving cytokine-mediated toxicity comprising administering to an individual an effective amount of an agent capable of inhibiting MIF receptor gene expression.  
25

17. The method of Claim 16 in which the agent is an antisense molecule complementary to MIF receptor mRNA.

18. The method of Claim 16 in which the agent is  
30 a ribozyme molecule specific for MIF receptor mRNA.

19. The method of Claim 16 in which the agent is a triple helix component.  
35

20. A method for treating a condition involving cytokine-mediated toxicity comprising administering to an individual an effective amount of an agent that inhibits MIF release.

5

21. The method of Claim 20 in which the agent inhibits steroid induced release of macrophage MIF.

22. The method of Claim 21 in which the agent is  
10 administered in combination with a steroid.

23. The method of Claim 20 in which the agent inhibits toxin induced release of macrophage MIF.

15 24. The method of Claim 20 in which the agent inhibits  $\text{TNF}\alpha$ -induced release of macrophage MIF.

25. The method of Claim 20 in which the agent inhibits  $\text{IFN-}\gamma$  induced release of macrophage MIF.

20

26. The method of Claim 20 in which the agent inhibits LPS induced release of pituitary MIF.

27. A method for treating a condition involving  
25 cytokine-mediated toxicity comprising administering to an individual an effective amount of (a) an agent capable of inhibiting MIF biologic activity, MIF receptor biologic activity, MIF gene expression, MIF receptor gene expression, or MIF release in  
30 combination with (b) anti- $\text{TNF}\alpha$ , anti-IL-1, anti- $\text{IFN-}\gamma$ , IL-1RA, a steroid, a glucocorticoid, or IL-10.

28. The method of Claim 27 in which the agent is an MIF-antagonist.

35

29. The method of Claim 28 in which the antagonist is an anti-MIF antibody.

30. The method of Claim 28 in which the  
5 antagonist is a soluble MIF-receptor.

31. The method of Claim 28 in which the MIF-antagonist is administered in combination with a soluble receptor for  $\text{TNF}\alpha$ , IL-1 or  $\text{IFN-}\gamma$ .

10

32. The method of Claim 28 in which the antagonist is a small organic molecule.

33. The method of Claim 1, 6, 11, 16, 20 or 27  
15 in which the condition is septic shock.

34. The method of Claim 1, 6, 11, 16, 20 or 27 in which the condition is inflammation.

20 35. The method of Claim 1, 6, 11, 16, 20 or 27 in which the condition is autoimmunity.

36. The method of Claim 1, 6, 11, 16, 20 or 27 in which the condition is cachexia.

25

37. The method of Claim 1, 6, 11, 16, 20 or 27 in which the condition is a viral infection.

38. The method of Claim 1, 6, 11, 16, 20 or 27  
30 in which the condition is a bacterial infection.

39. A method for inhibiting the toxic side effects of therapeutic steroids, comprising administering to an individual an effective amount of  
35 (a) an agent that modulates MIF biologic activity, MIF

receptor biologic activity, MIF gene expression, MIF receptor gene expression, or MIF release, and (b) the therapeutic steroid.

5           40. The method of Claim 39 in which the modulation is inhibition.

          41. A method for enhancing the anti-inflammatory activity of therapeutic steroids, comprising  
10 administering an effective amount of (a) an agent that modulates MIF biologic activity, MIF receptor biologic activity, MIF gene expression, MIF receptor gene expression, or MIF release, and (b) the therapeutic steroid.

15

          42. The method of Claim 41 in which the modulation is inhibition.

          43. A method for detecting levels of MIF in a  
20 body fluid as an indicator of a disease condition comprising:

- (a) contacting a test sample with an MIF binding partner under conditions and for a time sufficient to form MIF-binding partner complexes; and  
25
- (b) detecting the MIF-binding partner complexes.

          44. The method of Claim 43 in which the MIF  
30 binding partner is an anti-MIF antibody.

          45. A method for identifying compounds that inhibit cellular release of MIF, comprising:

35

- 5 (a) adding a test compound to a culture of cells that release MIF upon exposure to an MIF-inducer;
- (b) adding the MIF-inducer to the cell culture;
- (c) detecting any MIF released into the cell culture medium.

10 46. The method of Claim 45 in which the cells are a macrophage cell line.

47. The method of Claim 46 in which the MIF-inducer is a steroid.

15 48. The method of Claim 46 in which the MIF-inducer is a toxin.

49. The method of Claim 46 in which the MIF-inducer is  $\text{TNF}\alpha$ .

20

50. The method of Claim 46 in which the MIF-inducer is  $\text{IFN-}\alpha$ .

25 51. The method of Claim 45 in which the cells are a pituitary cell line.

52. The method of Claim 51 in which the MIF-inducer is a steroid.

30 53. The method of Claim 51 in which the MIF-inducer is a toxin.

54. An MIF receptor having an amino acid sequence comprising: Ile-X-His-Asn-Thr-Val-Ala-Thr-  
35 Glu-Ile-(Ser)-(Gly)-Tyr-Asn-(Asn/Gly)-(Ala)-(Met).



55. An MIF receptor having an amino acid sequence comprising: Ala-Lys-Lys-Gly-Ala-Val-Gly-Gly-Ile.

5 56. A cell line capable of expressing an exogenous MIF coding sequence.

57. A transgenic animal in which MIF gene expression is modified.

10

58. The transgenic animal of Claim 57 in which the modified MIF gene expression is inhibited.

59. A transgenic animal in which MIF receptor  
15 gene expression is modified.

60. The transgenic animal of Claim 59 in which the modified MIF receptor gene expression is inhibited.

20

61. An antibody that immunospecifically binds to MIF.

62. The antibody of Claim 61 which is a  
25 monoclonal antibody.

30

35




FIG.1


FIG. 2

start  
↓  
CCATGCCTATGTTTCATCGTGAACACCAATGTTCCCCGCGCC  
TCCGTGCCAGAGGGGTTTCTGTGGAGCTCACCAGCAGCTGGCGCAGGCCACCGGCAAG  
CCCGCACAGTACATCGCAGTGCAGTGGTCCCGGACCAGCTCATGACTTTTAGCGGCACG  
AACGATCCCTGCGCCCTCTGCAGCCTGCACAGCATCGGCAAGATCGGTGGTGCCCAGAAC  
CGCAACTACAGTAAGCTGCTGTGTGGCCTGCTGTCCGATCGCCTGCACATCAGCCCGGAC  
CGCTCCTACAGCAAGCTGCTGTGCGGCCTGCTGGCCGAGCGCCTGCGCATCAGCCCGGAC  
CGGGTCTACATCAACTATTACGACATGAACGCTGCCAACGTGGGCTGGAACGGTTCCACC  
AGGGTCTACATCAACTATTACGACATGAACCGGCCAGTGTGGGCTGGAACAACCTCCACC  
TTCGCTTGAGTCCTGGCCCCACTTACCTGCACCGCTGTTC  
↑  
stop

FIG.3

	10	20	30	40	50	60
mu	MPMF	I VNTN	VPRA	SVPE	GF	SEL
						TQQL
						AQAT
						GKPA
						QYIA
						VHV
						PDQL
						MTF
						SGT
						NDP
						PCAL
						C
hu	MPMF	I VNTN	VPRA	SVPE	GF	SEL
						TQQL
						AQAT
						GKPP
						QYIA
						VHV
						PDQL
						MAF
						GGS
						SEPC
						ALC
	70	80	90	100	110	
mu	SLHS	IGG	AQNR	SYSK	LLCG	LLSD
						RLHI
						SPDR
						VYIN
						YYDM
						NAAN
						VGVN
						GS
						TFA
hu	SLHS	IGG	AQNR	SYSK	LLCG	LLAE
						RLRI
						SPDR
						VYIN
						YYDM
						NAAN
						VGVN
						GS
						TFA

FIG.4

**FIG. 5**

5/22




FIG. 6A




FIG. 6B

6/22

RECTIFIED SHEET (RULE 91)




FIG.7

7/22

RECTIFIED SHEET (RULE 91)






FIG. 8




FIG. 9B




FIG. 9A




FIG. 10




FIG. 11




FIG. 12




FIG.13



FIG. 14B




FIG. 14A








FIG. 15





**FIG. 17**





FIG. 18

**FIG. 19**




FIG. 20





FIG. 21




FIG. 22

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/05433**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : A61K 39/395; C07K 15/28; C12N 5/12

US CL : 424/130.1, 133.1, 134.1, 143.1

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1, 133.1, 134.1, 143.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG, BIOSIS, EMBASE, MEDLINE, CA, WPI

search terms: migration inhibitory factor, mif, sepsis, inflammation, bucala

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. IMMUNOL., VOLUME 137, ISSUED 15 JULY 1986, D.Y. LIU ET AL., "A MONOCLONAL ANTIBODY SPECIFIC FOR A MONOCYTE-MACROPHAGE MEMBRANE COMPONENT BLOCKS THE HUMAN MONOCYTE RESPONSE TO MIGRATION INHIBITORY FACTOR", PAGES 448-455, SEE ENTIRE DOCUMENT.	1-3, 33-38
Y	PROC. NATL. ACAD. SCI., VOLUME 86, ISSUED OCTOBER 1989, W.Y. WEISER ET AL., "MOLECULAR CLONING OF A cDNA ENCODING A HUMAN MACROPHAGE MIGRATION INHIBITORY FACTOR", PAGES 7522-7526, SEE ENTIRE DOCUMENT.	1-3, 33-38

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

19 AUGUST 1994

Date of mailing of the international search report

AUG 29 1994

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

PHILLIP GAMBEL, Ph.D.

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/05433

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. INFECT. DIS., VOLUME 163, ISSUED 1991, C.A. DINARELLO , "THE PROINFLAMMATORY CYTOKINES INTERLEUKIN-1 AND TUMOR NECROSIS FACTOR AND TREATMENT OF THE SEPTIC SHOCK SYNDROME", PAGES 1177-1184, SEE ENTIRE DOCUMENT.	1-3, 33-38
X,P	NATURE, VOLUME 365, ISSUED 21 OCTOBER 1993, J. BERNHAGEN ET AL., "MIF IS A PITUITARY-DERIVED CYTOKINE THAT POTENTIATES LETHAL ENDOTOXAEMIA", PAGES 756-759, SEE ENTIRE DOCUMENT.	1-3, 33-38

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING**

This ISA found multiple inventions as follows:

- I. Claims 1-3, and 33-38, drawn to a method wherein an anti-MIF antibody is administered to antagonize MIF activity *in vivo*, classified in Class 424, subclass 85.91, for example.
- II. Claims 1, 2, 4, and 33-38, drawn to a method wherein a soluble MIF receptor is administered to antagonize MIF activity *in vivo*, classified in Class 514, subclass 2, for example.
- III. Claims 1, 2, 5, and 33-38, drawn to a method wherein a small organic molecule is administered to antagonize MIF activity *in vivo*, classified in Class 514, subclass 1, for example.
- IV. Claims 6-8, and 33-38, drawn to a method wherein an anti-MIF-receptor (anti-MIF-R) antibody is administered to inhibit MIF-R activity *in vivo*, classified in Class 424, subclass 85.8, for example.
- V. Claims 6, 7, 9, and 33-38, drawn to a method wherein a MIF analog is administered to inhibit MIF-R activity *in vivo*, classified in Class 514, subclass 2, for example.
- VI. Claims 6, 7, 10, and 33-38, drawn to a method wherein a small organic molecule is administered to inhibit MIF-R activity *in vivo*, classified in Class 514, subclass 1, for example.
- VII. Claims 11, 12, and 33-38, drawn to a method wherein an antisense oligonucleotide complementary to MIF mRNA is administered to inhibit MIF gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- VIII. Claims 11, 13, and 33-38, drawn to a method wherein a ribozyme specific for MIF mRNA is administered to inhibit MIF gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- IX. Claims 11, 14, and 33-38, drawn to a method wherein a triplex-forming oligo specific for MIF mRNA is administered to inhibit MIF gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- X. Claims 11, 15, and 33-38, drawn to a method wherein a steroid is administered to inhibit MIF gene expression *in vivo*, classified in Class 514, subclass 169, for example.
- XI. Claims 16, 17, and 33-38, drawn to a method wherein an antisense oligo complementary to MIF-R mRNA is administered to inhibit MIF-R gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- XII. Claims 16, 18, and 33-38, drawn to a method wherein a ribozyme specific for MIF-R mRNA is administered to inhibit MIF-R gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- XIII. Claims 16, 19, and 33-38, drawn to a method wherein a triplex-forming oligo specific for MIF-R mRNA is administered to inhibit MIF-R gene expression *in vivo*, classified in Class 514, subclass 44, for example.
- XIV. Claims 20-26 and 33-38, drawn to a method wherein an agent is administered to inhibit MIF release *in vivo*, classified in Class 514, subclass 789, for example.
- XV. Claims 27-29 and 33-42, drawn to a method wherein an anti-MIF antibody is co-administered with an anti-cytokine antibody, or a steroid, or IL-10, classified in Class 424, subclass 85.91, for example.
- XVI. Claims 27, 28, 30, and 33-42, drawn to a method wherein a soluble MIF receptor is co-administered with an anti-cytokine antibody, or a steroid, or IL-10, classified in Class 514, subclass 2, for example.
- XVII. Claims 27, 28, 31, and 33-42, drawn to a method wherein a MIF antagonist is co-administered with a soluble receptor for a different cytokine, classified in Class 514, subclass 2, for example.
- XVIII. Claims 27, 28, and 32-42, drawn to a method wherein a small organic molecule which inhibits activity of MIF is co-administered with an anti-cytokine antibody, or a steroid, or IL-10, classified in Class 514, subclass 171, for example.
- XIX. Claims 43-53, drawn to a method for detecting levels of MIF in a body fluid using anti-MIF antibodies, and to a method for identifying compounds which alter extracellular levels of MIF, classified in Class 436,

subclass 536, for example.

- XX. Claims 54 and 55, drawn to MIF receptor protein, classified in Class 530, subclass 350, for example.
- XXI. Claim 56, drawn to a cell line that expresses an exogenous MIF gene, classified in Class 435, subclass 240.2, for example.
- XXII. Claims 57 and 58, drawn to a transgenic animal in which MIF gene expression is modified, classified in Class 800, subclass 2, for example.
- XXIII. Claims 59 and 60, drawn to a transgenic animal in which MIF receptor gene expression is modified, classified in Class 800, subclass 2, for example.
- XXIV. Claims 61 and 62, drawn to antibodies that bind MIF, classified in Class 530, subclass 387.2, for example.

This application contains the above-listed groups of separate inventions which are not linked so as to form a single general inventive concept as required by PCT Rule 13.1. The technical features which link Group I to each of the other Groups II-XXIV are not advances over the prior art within the meaning of PCT Rule 13.2.

The method of Group I wherein an anti-MIF antibody is administered, and the different methods of Groups II and XVI, wherein a soluble MIF receptor is administered, have in common the special technical feature that the agent administered is a lymphokine inhibitor; however the special technical feature of the inhibition of a lymphokine by administration of a soluble receptor which links Group I to Group II and Group XVI is not an advance over the prior art as required by PCT Rule 13.2. Vassalli teaches injection of soluble receptor to inhibit activity of the lymphokine which binds the receptor (Annual Review of Immunology, Volume 10 (1992) pp. 411-452, see p. 430).

The method of Group I wherein an anti-MIF antibody is administered, and the different methods of Groups III and XVIII, wherein a small organic molecule is administered, have in common the special technical feature that the agent administered inhibits lymphokine activity; however the special technical feature of the inhibition of a lymphokine by administration of a small organic molecule which links Group I to Group III and Group XVIII is not an advance over the prior art as required by PCT Rule 13.2. Sigal et al. teach injection of small organic molecules to inhibit lymphokine-mediated events (Annual Review of Immunology, Volume 10 (1992) pp. 519-560, see pp. 520-525).

The method of Group I wherein an anti-lymphokine antibody is administered, and the methods of Groups IV, V, and VI, wherein a MIF-R antagonist is administered, have in common the special technical feature that the diverse agents recited in claims IV, V, and VI all inhibit lymphokine activity; however the special technical feature of the inhibition of a lymphokine by administration of an antagonist which binds and inhibits the lymphokine receptor, which links Group I to Groups IV, V, and VI, is not an advance over the prior art as required by PCT Rule 13.2. Vassalli (p. 426) and Smith (Science, Volume 240 (27 May 1988) pp. 1169-1176, see p. 1175) both teach that the activity of a lymphokine can be inhibited by an antagonist of the lymphokine receptor.

The method of Group I wherein an anti-lymphokine antibody is administered, and the methods of Groups VII, VIII, and IX, wherein an oligonucleotide with antisense, ribozyme, or triplex-forming activity, respectively, is administered, have in common the special technical feature that the diverse inhibitors of lymphokine gene expression recited in claims VII, VIII, and IX, all inhibit lymphokine activity; however the special technical feature of the inhibition of activity of a given protein by administration of an oligo which inhibits expression of the gene encoding said protein, which links Group I to Groups VII, VIII, and IX, is not an advance over the prior art as required by PCT Rule 13.2. Uhlmann et al. teach the methods whereby expression of a given gene is inhibited by administration of oligonucleotides with antisense, ribozyme, or triplex-forming activity (Chemical Reviews, Volume 90, No. 4 (June 1990) pp. 543-584, see pp. 574-579).

The method of Group I wherein an anti-lymphokine antibody is administered, and the method of Group X, wherein a steroid is administered, have in common the special technical feature that the agent administered inhibits lymphokine activity; however the special technical feature of the inhibition of a lymphokine by administration of a steroid which inhibits expression of the lymphokine gene, which links Group I to Group X, is not an advance over the prior art as required by PCT Rule 13.2. Smith teaches a method wherein expression of a lymphokine gene is inhibited by administration of a steroid (p. 1175).

The method of Group I wherein an anti-lymphokine antibody is administered, and the methods of Groups XI, XII, and XIII, wherein an oligonucleotide with antisense, ribozyme, or triplex-forming activity, respectively, is administered, have in common the special technical feature that the diverse agents recited in claims XI, XII, and XIII, all inhibit

lymphokine activity by inhibiting expression of the lymphokine receptor gene; however the special technical feature of the inhibition of a lymphokine by administration of an oligo which inhibits expression of the lymphokine receptor gene, which links Group I to Groups XI, XII, and XIII, is not an advance over the prior art as required by PCT Rule 13.2. Uhlmann et al. teach the method wherein expression of a given gene is inhibited by administration of oligonucleotide with antisense, ribozyme, or triplex-forming activity (pp. 574-579), and Vassalli (p. 426) and Smith (p. 1175) both teach that the activity of a lymphokine can be inhibited by an agent which inhibits activity of the lymphokine receptor, as noted above.

The method of Group I wherein an anti-lymphokine antibody is administered, and the method of Group XIV, wherein an inhibitor of lymphokine release is administered, have in common the special technical feature that the agent administered inhibits lymphokine activity; however the special technical feature of the inhibition of activity of a lymphokine by administration of an agent which inhibits lymphokine release, which links Group I to Group XIV, is not an advance over the prior art as required by PCT Rule 13.2. Smith teaches a method wherein activity of a lymphokine is inhibited by administration of an agent which prevents lymphokine release (p. 426).

The method of Group I wherein an anti-lymphokine antibody is administered, and the different methods of Groups XV, XVI, and XVII, wherein an antagonist to a different lymphokine is co-administered, have in common the special technical feature that the agents are co-administered to inhibit lymphokine activity and modulate an immune response; however the special technical feature of co-administration of an anti-lymphokine antibody and an antagonist of a different lymphokine to modulate an immune response, which links Group I to Groups XV, XVI, or XVII, is not an advance over the prior art as required by PCT Rule 13.2. Vassalli teaches a method wherein antibodies that bind and inhibit one lymphokine are co-administered with an antagonist of a different lymphokine to modulate an immune response (pp. 430).

The method of Group I wherein an anti-lymphokine antibody is administered, and the method of Group XIX wherein an anti-lymphokine antibody is used to detect levels of the lymphokine in a biological sample, have in common the special technical feature that an antibody specific for the lymphokine is used; however the use of an anti-lymphokine antibody to detect levels of the lymphokine in a biological sample, which links Group I to Group XIX, is not an advance over the prior art as required by PCT Rule 13.2. Peck teaches a method wherein an anti-lymphokine antibody is used to detect levels of the lymphokine in a biological sample (Journal of Immunological Methods, Volume 64 (1983) pp. 179-187).

The method of Group I wherein an anti-lymphokine antibody is administered, and the invention of Group XX, which is the sequenced lymphokine receptor protein, have in common the special technical feature that the lymphokine binds specifically to the receptor; however the special technical feature of the specific relationship between a lymphokine and its sequenced receptor, which links Group I to Group XX, is not an advance over the prior art as required by PCT Rule 13.2. Holmes et al. disclose the amino acid sequence of a receptor that specifically binds a lymphokine (Science, Volume 253 (13 September 1991) pp. 1278-1280).

The method of Group I wherein an anti-lymphokine antibody is administered, and the cell line of Group XXI which expresses an exogenous gene, have in common the special technical feature that the exogenous gene encodes the lymphokine of Group I; however the special technical feature of the expression of an exogenous lymphokine gene in a cell line, which links Group I to Group XXI, is not an advance over the prior art as required by PCT Rule 13.2. Weiser et al. disclose a transformed cell line which expresses an exogenous lymphokine gene (Proceedings of the National Academy of Sciences, Volume 86 (October 1989) pp. 7522-7526, see pp. 7522-7523).

The method of Group I wherein an anti-lymphokine antibody is administered, and the transgenic animal of Group XXII, have in common the special technical feature of the modified expression of the lymphokine gene in a transgenic animal; however the special technical feature of the modified expression of a lymphokine gene in a transgenic animal, which links Group I to Group XXII, is not an advance over the prior art as required by PCT Rule 13.2. Buhler et al. disclose a transgenic animal wherein an exogenous lymphokine gene is expressed (Bio/Technology, Volume 8 (February 1990) pp. 140-143).

The method of Group I wherein an anti-lymphokine antibody is administered, and the transgenic animal of Group XXIII, have in common the special technical feature of the modified expression of a gene encoding a lymphokine receptor in a transgenic animal; however the special technical feature of the modified expression of a gene encoding a lymphocyte surface receptor for an immuno modulator in a transgenic animal, which links Group I to Group XXIII, is not an advance over the prior art as required by PCT Rule 13.2. Fung-Leung et al. disclose a transgenic animal wherein expression in lymphocytes of an exogenous gene encoding a surface-bound, immunomodulating receptor is inhibited (Cell, Volume 65 (03 May 1991) pp. 443-449).

The method of Group I wherein an anti-lymphokine antibody is administered, and the anti-lymphokine antibodies of

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/05433

Group XXIV, have in common the special technical feature that the antibodies of Group XXIV are used to inhibit lymphokine activity; however the special technical feature of the inhibition of a lymphokine by administration of the anti-lymphokine antibodies, which links Group I to Group XXIV, is not an advance over the prior art as required by PCT Rule 13.2. Vassalli teaches a method wherein anti-lymphokine antibodies are administered to inhibit lymphokine activity (p. 430).